



Marengo®

Product information sheet

- // Active Ingredient: Indaziflam 7.4%
- // IRAC Group: 29
- // EPA Registration Number: 432-1518
- // Signal Word: None
- // Formulation: Soluble Concentrate
- // REI: 12 hours

Marengo® preemergence herbicide provides long-lasting weed control on over 70 weeds.

Best Uses


- // Broadcast or directed spray in nurseries and non-crop settings to selectively prevent broadleaf weeds, grasses and annual sedges
- // Apply in fall for continuous weed control into the following growing season
- // Effective on hard-to-control weeds such as eclipta, bittercress, groundsel, oxalis, marestail, spurge, crabgrass and annual bluegrass
- // The only preemergence herbicide labeled for use inside greenhouses beneath benches

Attributes

- // Up to 8 months of residual control
- // Reduces labor costs with fewer applications and less clean up work
- // Labeled for use on 150 ornamental plants
- // Indaziflam has a unique mode of action making it ideal for rotation and resistance management
- // No signal word
- // Irrigation or rainfall is recommended within 21 days of application


Application Rates

- // For broadcast applications, use 7.5-15.5 oz./A
- // For directed spray applications, use 0.17-0.36 fl. oz./1,000 sq. ft.
- // Marengo is available in 18-oz. and 64-oz. containers



Preemergence Weed Control With Marengo®

3 Months After Application



backedbybayer.com | backedbybayer.com/production-ornamentals



ALWAYS READ AND FOLLOW LABEL INSTRUCTIONS
Bayer CropScience LP, Environmental Science Division, 2 T.W. Alexander Drive, Research Triangle Park, NC 27709. For additional product information, call toll-free 1-800-331-2867. Not all products are registered in all states. Bayer, the Bayer Cross, and Marengo are registered trademarks of Bayer. Gallery is a registered trademark of Dow. Sureguard is a registered trademark of Valent.
© 2018 Bayer CropScience LP. ES-318-MAR-71-A-R1