



FOR SUPERIOR CONTROL OF KEY DISEASES IN TURFGRASS AND ORNAMENTALS

- Multi-site
- Very low chance of resistance buildup
- Economical
- Broad-spectrum
- Great tank-mix partner

FRAC class: M3

Active ingredient: *Mancozeb*

Formulation: WDG | Liquid

Package size: 8 x 6 lb | 2 x 2.5 gallon

TURFGRASS & SODFARMS Diseases Controlled

Algae	Pink Snow Mold
Brown Patch	Pythium Blight
Copper Spot	Red Thread
Dollar Spot	Rhizoctonia Solani
Fusarium Blight	Slime Molds
Gray Leaf Spot	Stem Rust
Leaf Rust	Stripe Rust
Leaf Spot	

Resistance Management Tool

Managing fungicide resistance is an important decision when choosing your disease control products. The multi-site action of Manzate T&O fungicide prevents pathogenic organisms from adapting and mutating, meaning even after decades of use, fungal strains have not developed resistance to Manzate T&O formulations.

Formulation	Use Rates*	Application Timing
Manzate Pro-Stick T&O	4 - 8 oz per 1,000 sq. ft.	Typically begin application when disease threatens, or at the first sign of disease. Repeat application as per labeled interval rates as conditions persist.
Manzate Max T&O	5 - 14 fl oz per 1,000 sq. ft.	

*Follow label instructions for disease specific use rates, application timing, and complete directions for use.





FLOWERS, FOLIAGE PLANTS AND ORNAMENTALS

Manzate T&O fungicide is recommended for use on certain flower, foliage, and ornamental plants. Please note the use of Manzate T&O has been found to be acceptable in specific genera and species of the plants listed on the product label; however, phytotoxicity may occur. Due to the large number of species and

varieties of ornamentals and nursery plants, each one is not tested for sensitivity to Manzate T&O. The user should determine if Manzate T&O can be used safely prior to commercial use. Manzate T&O may also be used in aerial applications, as per label specifications. See label for full listing of plants.

Diseases Controlled

Alternaria	Curvularia	Gloeosporium	Melampsoridium	Ramularia
Asochyta	Cylindrosporium	Glomerella	Microsphaera alni	Rhizoctonia
Bipolaris	Diaporthe	Gnomonia	Monilinia	Rhytisma
Botryosphaeria	Didymascella	Guignardia	Monochaetia	Sclerotium
Botrytis	Diplocarpon	Gymnosporangium	Mycosphaerella	Sirococcus
Cephalosporium	Drechslera	Helminthosporium	Ovulinia	Sphaeropsis
Cercospora	Elsinoe	Kabatiella	Peronospora	Stemphylium
Ciborinia	Entomosporium	Keithia	Pestalotia	Stigmia
Cladosporium	Erwinia	Leptosphaeria	Phaeocryptopus	Taphrina
Coccomyces	Exobasidium	Lophodermium	Phoma	Uromyces
Colletotrichum	Fabraea	Marssonina	Phomopsis	Venturia
Coryneum	Fusarium	Melampsora	Phyllosticta	Whetzelinia
	Fusicladium		Puccinia	

Formulation

Use Rates*

Application Timing

Manzate Pro-Stick T&O	1 - 2 lbs. per acre or 100 gals. water	Typically begin application when disease threatens, or at the first sign of disease. Repeat application as per labeled interval rates as conditions persist.
Manzate Max T&O	0.8 - 1.6 quarts per acre or 100 gals. water	

*Follow label instructions for disease specific use rates, application timing, and complete directions for use.



Contact your local UPI sales representative or distributor, call 1-800-438-6071 or visit www.upi-usa.com to learn more.

Always read and follow label directions and precautions. Manzate® and UPI logo are registered trademarks of United Phosphorus, Inc. ©June, 2017. UPI, 630 Freedom Business Center, King of Prussia, PA 19406.

MANZATE-SPD-17-01