



HERBICIDE

# 2,4-D LV 4

2,4-D LV 4 is a low-volatile phenoxy herbicide formulation that controls a variety of broadleaf weeds in crops including cereal grains, corn, sorghum, soybeans (pre-plant burndown only), rangeland & pasture, fallow land and ornamental turf. 2,4-D LV 4 will be more effective at controlling weeds at temperatures below 50°F as compared to 2,4-D amine formulations.

## PRODUCT FEATURES:

- A low-volatile formulation that has excellent compatibility with a variety of fertilizer solutions.
- Effective in controlling more than 90 broadleaf weeds when applied at the young and active growing stage.
- 2,4-D LV 4 offers broad label uses including forestry, grasses grown for turf, red potatoes and various non-crop applications.

## APPLICATION RESTRICTIONS

- 2,4-D esters may volatilize during conditions of low humidity and high temperatures. Do not apply 2,4-D LV 4 during these environmental conditions.
- Higher use rates of 2,4-D LV 4 may increase the risk of crop injury and should be used only where weed control justifies such a risk.
- Applicators must follow all state and local pesticide drift requirements regarding application of 2,4-D herbicides.
- **2,4-D LV 4 is NOT approved for application to the new Corn, Cotton and Soybean 2,4-D tolerant GMO crops.**

See label for complete application guidelines and restrictions.

ACTIVE INGREDIENT: 2,4-D ESTER

HERBICIDE GROUP: I O

ACTIVE INGREDIENT PER GALLON: 3.8 lbs.

MODE OF ACTION: Synthetic auxin growth regulator

USE RATES: See label for all product use rates

PACKAGING: 4x1's, 2x2.5's, 30's and 265 gal



Albaugh®  
Your Alternative™

ALBAUGHLLC.COM





## WEEDS CONTROLLED

### ANNUAL OR BIENNIAL WEEDS

Beggarticks <sup>(1)</sup>	Kochia	Radish, Wild
Bittercress,	Lamsquarter, Common	Ragweed, Common
Smallflowered <sup>(2)</sup>	Lettuce, Prickly <sup>(1)(2)</sup>	Ragweed, Giant
Bitterweed	Lettuce, Wild	Rape, Wild
Broomweed, Common <sup>(1)</sup>	Lupines	Rocket, Yellow
Burdock, Common	Mallow, Little <sup>(1)</sup>	Salsify, Common <sup>(1)</sup>
Buttercup,	Mallow, Venice <sup>(1)</sup>	Salsify, Western <sup>(1)</sup>
smallflowered <sup>(1)(2)</sup>	Marshelder	Shepherdspurse
Carpetweed	Morningglory, Annual	Sicklepod
Cinquefoil, Common <sup>(2)</sup>	Morningglory, Ivy	Smartweed
Cinquefoil, Rough <sup>(2)</sup>	Morningglory, Woolly	(Annual Species) <sup>(1)(2)</sup>
Cocklebur, Common	Mousetail <sup>(2)</sup>	Sneezeweed, Bitter
Coffeeweed	Mustards (except	Sowthistle, Annual
Copperleaf, Virginia <sup>(2)</sup>	blue mustard)	Sowthistle, Spiny
Croton, Texas	Parsnip, Wild	Spanishneedles
Croton, Wooly	Pennycress (Fanweed)	Sunflower
Fixweed	Pepperweeds	Sweetclover
Galinsoga	( <i>Lepidium</i> Spp.) <sup>(1)(2)</sup>	Tansymustard
Geranium, Carolina <sup>(2)</sup>	Pigweeds	Thistle, Bull
Hemp, Wild	( <i>Amaranthus</i> Spp.) <sup>(1)</sup>	Thistle, Musk <sup>(1)</sup>
Horseweed(Marestail) <sup>(2)</sup>	<i>Poorjoe</i>	Thistle, Russian
Jewelweed	Primrose, Common	(Tumbleweed) <sup>(1)</sup>
Jimsonweed	Purslane, Common <sup>(2)</sup>	Velvetleaf
Knotweed <sup>(1)</sup>	Pusley, Florida	Vetches

### PERENNIAL WEEDS

Alfalfa <sup>(1)(2)</sup>	Chicory	Ironweed, Western <sup>(2)</sup>
Austrian Fieldcress <sup>(1)</sup>	Clover, Red <sup>(1)(2)</sup>	Ivy, Ground <sup>(1)</sup>
Artichoke, Jerusalem <sup>(1)</sup>	Coffeeweed	Nettles (Including Stinging) <sup>(1)</sup>
Aster, many-flower <sup>(1)</sup>	Cress, Hoary <sup>(1)</sup>	Onion, Wild <sup>(1)</sup>
Bindweed (hedge, field	Dandelion	Pennywort
and European) <sup>(1)(2)</sup>	Docks <sup>(1)</sup>	Plantains
Blue Lettuce	Dogbanes <sup>(1)</sup>	Ragwort, Tansy <sup>(1)</sup>
Blueweed, Texas	Eveningprimrose, Cutleaf <sup>(2)</sup>	Sowthistle, Perennial
Broomweed	Garlic, Wild	Thistle, Canada <sup>(1)(2)</sup>
Bullnettle <sup>(1)(2)</sup>	Goldenrod	Vervains <sup>(1)</sup>
Carrot, Wild <sup>(1)</sup>	Hawkweed, Orange <sup>(1)</sup>	Wormwood
Catnip	Healall	

<sup>(1)</sup> Difficult-to-Control Weeds: These weeds are only partially controlled and may require repeat applications and/or use of the higher specified rate of this product even under ideal conditions of application.

<sup>(2)</sup> This product may not be used to control this weed species in the state of California.

## PRODUCT USE RATES

### PREPLANT/PRE-EMERGENT APPLICATION

Corn	1 to 2 pints/A
Soybeans*	¾ to 2 pints/A

### POST-EMERGENT APPLICATION

Cereal Grains	½ to 2 pints/A
Corn	½ to 1 pint/A
Ornamental Turf	¾ to 3 pints/A
Sorghum	½ to 1 pint/A
Rangeland/Fallow	1 to 4 pints/A

\*Not registered for use in California

Always read and follow label directions.  
EPA Reg. No. 42750-15 | AD100410