

Patriot® Selective Herbicide controls tough broadleaf weeds such as Russian thistle and mustards in wheat, barley, fallow, pastures and rangeland. It provides an economical post-patent alternative. With contact and residual control, Patriot burns down existing weeds and keeps late flushes down for four to six weeks. And its wide window of application gives you flexibility if application is delayed.

COMPETITIVE ADVANTAGES

- Excellent post-emerge weed control
- · Offers excellent crop safety
- Provides residual control for up to six weeks
- Wide application window
- Excellent tank-mix partner

AT-A-GLANCE INFO

EPA REG. NO. 228-391

ACTIVE INGR. metsulfuron-methyl (60%)

FORMULATION water dispersible

granule

CHEM. FAMILY sulfonylurea

HRAC NO.

MODE OF ACT. ALS inhibitor

4 hours RFI

SIGNAL WORD caution

RESTRICTED

PACKAGE SIZE 16 oz bag (8 per case)













COUNT ON PATRIOT® FOR BROADLEAF WEED CONTROL.

KEY USES

Barley

Conservation reserve program (CRP) land

Fallow

Grain sorghum

Pastures

Rangeland

Triticale

Wheat

KEY WEEDS CONTROLLED

(See product label for complete list)

Annual marshelder

Bitter sneezeweed

Blackeyed-Susan

Broomweed, common

Bur buttercup

Carolina geranium

Chamomile; false, mayweed

Chickweed, common

Coast fiddleneck (tarweed)

Conical catchfly

Cowcockle

Curly dock

Dandelion

Dogfennel

Field pennycress (fanweed)

Filaree

Flixweed

Groundsel

Henbit

Kochia

Lambsquarters

Lettuce; miners, prickley

Marestail

Mustard; tansy, treacle, tumble, wild

Pigweeds

Plains coreopsis

Purslane, common

Shepherd's-purse

Smallseed falseflax

Smartweeds

Sunflower, volunteer

Thistle, Russian

Waterpod

Woolly croton

Yarrow

USE SUMMARY

APPLICATION TIMING AND RATES

Wheat and Barley: Apply 1/10 oz/A. To avoid crop injury, do not apply during boot or early heading. For dryland wheat and barley (except durum or wampum variety), apply after the crop is in the 2-leaf stage, but before boot. For durum and wampum variety spring wheat, make applications after tillering, but before boot. Applications to durum and wampum varieties should be made in combination with 2,4-D. For irrigated wheat and barley, apply after the crop begins tillering, but before boot. First post-treatment irrigation should be delayed for at least 3 days after treatment and should not exceed one inch of water.

Pasture and Rangeland: Apply 1/10 to 4/10 oz/A as a broadcast spray. For spot treatments, use up to 1 oz per 100 gal of water.

Harvest Aid: Apply 1/10 oz/A in combination with 2,4-D (or Credit® 41 Extra/Credit® Xtreme/Razor® Pro) to aid in dry down of many broadleaf weeds. Make applications after the crop has reached the hard dough stage, but no later than 10 days before harvest.

Fallow: Patriot® may be used as a fallow treatment in the spring or fall when the majority of weeds have emerged and are actively growing. Apply at 1/10 oz/A.

Grain Sorghum: Apply 1/20 oz/A of Patriot plus 1/4 lb active ingredient of 2,4-D amine per acre. Do not use a surfactant or crop oil.

See label for complete directions for use.



