

Supplemental Label

Poast[®]

Herbicide

For control of annual and perennial grass weeds in fine fescue grown for turfgrass seed in Idaho, Oregon, and Washington Only

This supplemental label expires June 1, 2019, and must not be used or distributed after this date.

Active Ingredient*:

sethoxydim: 2-[1-(ethoxyimino)butyl]-5-[2-(ethylthio)propyl]-3-hydroxy-2-cyclohexen-1-one 18.0%

Other Ingredients: 82.0%

Total: 100.0%

*Equivalent to 1.5 pounds of sethoxydim per gallon

Contains petroleum distillate

EPA Reg. No. 7969-58

WARNING/AVISO

Directions For Use

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- The supplemental labeling and the entire **Poast[®] herbicide** container label, EPA Reg. No. 7969-58, must be in possession of the user at the time of application.
- Read the label affixed to the container for **Poast** before applying.
- Use of **Poast** according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for **Poast**.

Product Information

Poast is a broad-spectrum, postemergence herbicide for selective control of annual and perennial grass weeds. Avoid all direct or indirect contact with any desired grass crop (e.g., corn, rice, small grains, sorghum, ornamental grasses, and turfgrass).

Poast does not control sedges or broadleaf weeds.

Refer to the **Poast** container label for a complete list of weeds controlled.

Application Instructions

For use only in the states of Idaho, Oregon, and Washington

- **PHI** - 14 days
- **Maximum Single Application Rate** - 2.5 pints/acre
- **Maximum Seasonal Application Rate** - 5.0 pints/acre
- **DO NOT** apply more than 2 applications per season. Allow 30 days between applications.
- There are no livestock grazing or feeding restrictions in fine fescue grown for turfgrass seed.
NOTE: Treated fescue screenings or hay may be fed to animals.
- Ground application allowed. **DO NOT** apply by air.

Only apply **Poast** to fine fescue varieties classified as creeping red (*Festuca rubra*), Chewings (*Festuca nigrescens*), and hard fescue (*Festuca ovina*). Use standard high-pressure pesticide hollow cone or flat fan nozzles (minimum of 40 PSI at the nozzle); apply at least 10 gallons of water per acre. If foliage is dense, increase water volume to 20 gallons/acre (minimum of 60 PSI). Thorough foliar spray coverage is essential.

- **DO NOT** apply to tall fescue (*Schedonorus arundinaceus*) or any other desired grass plant. Injury will occur.

- **Poast® herbicide** does **NOT** control annual bluegrass (*Poa annua*) or rattail fescue (*Vulpia myuros*).
- **DO NOT** apply if rainfall is expected within 1 hour of application.
- **DO NOT** use flood or whirl chamber nozzles or selective application equipment (e.g., recirculating sprayers, wiper applicators, shielded applicators).

For control of annual ryegrass, colonial and highland bentgrasses, cheatgrass/downy brome, and German velvetgrass, apply when fine fescue is semi-dormant (generally November 1 through March 15). Application at other times of the year can result in reduced control. Refer to **Table 1** for application rate and timing (maximum height) information.

Table 1. Application Rate and Timing (Maximum Height) - Fine Fescue Grown for Turfgrass Seed		
Turfgrass Species	Maximum Height* (inches)	Rate/Acre** (pints)
Annual Turfgrass		
Annual ryegrass (<i>Lolium multiflorum</i>)	4 to 8	1.5
Cheatgrass/Downy brome (<i>Bromus tectorum</i>)	2 to 6	2.5
NOTE: Late fall applications after maximum germination produce the best results.		
Perennial Turfgrass		
German velvetgrass (<i>Holcus mollis</i>)	2 to 4	2.0 to 2.5
Colonial bentgrass Highland bentgrass (<i>Agrostis tenuis</i>)	2 to 4	1.5 to 2.5
NOTE: Use the higher rate on well-established grass weeds.		
*When fine fescue is semi-dormant		
**If regrowth occurs or new plants emerge, make a second application at the same rate.		

Conditions of Sale and Warranty

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND BASF'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF THE PRODUCT.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, EXEMPLARY, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF. 1108

Poast is a registered trademark of BASF.

© 2018 BASF Corporation
All rights reserved.

007969-00058.20180629.NVA 2018-04-025-0055
Supersedes: NVA 2013-04-025-0033

BASF Corporation
26 Davis Drive
Research Triangle Park, NC 27709



We create chemistry