

Supplemental Label

Pylon[®]
miticide-insecticide

This supplemental label expires on February 28, 2023, and must not be used or distributed after this date.

For use in commercial greenhouses on basil and chives sold for fresh use, and tomatoes less than 2 inches in diameter at maturity

Active Ingredient:

Chlorfenapyr: 4-bromo-2-(4-chlorophenyl)-1-(ethoxymethyl)- 5-(trifluoromethyl)-1*H*-pyrrole-3-carbonitrile..... 21.4%

Other Ingredients:..... 78.6%

Total: 100.0%

(8 fluid ounces contains 0.125 pounds of active ingredient.)

(1 gallon contains 2 pounds of active ingredient.)

EPA Reg. No. 241-374

Directions For Use

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- The supplemental labeling and the entire **Pylon**[®] miticide-insecticide container label, EPA Reg. No. 241-374, must be in possession of the user at the time of application.
- Read the label affixed to the container for **Pylon** before applying.
- Use of **Pylon** according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for **Pylon**.

Application Instructions

To limit the potential for resistance development, rotate the use of **Pylon** or other **Group 13** miticides/insecticides within a growing season, or among growing seasons, with different groups that control the same pests. Avoid application of more than the maximum seasonal use rate or the total number of consecutive sprays of **Pylon** per season.

Apply **Pylon** according to the rates, timing and resistance-management instructions in the **Application Rates and Instruction Remarks** table in this label.

Use Restrictions

Refer to the **Pylon** container label for specific use directions, recommendations and restrictions.

Application Rates¹ and Instruction Remarks

Crop	Target Pests	Rate (fl ozs of Pylon [®] miticide-insecticide/A)	Rate (lb ai/A)	Remarks
Greenhouse Crops: Basil and Chives sold for fresh use Tomatoes less than 2 inches in diameter when mature Cultivars, varieties and/or hybrids of these	Caterpillars, including: Beet armyworm <i>Spodoptera exigua</i> Southern armyworm <i>S. eridania</i> Fall armyworm <i>S. frugiperda</i> Yellowstriped armyworm <i>S. ornithogalli</i> Tomato pinworm <i>Keiferia lycopersicella</i> Tomato fruitworm <i>Heliothis zea</i> Hornworms <i>Manduca quinquemaculata</i> Cabbage looper <i>Trichoplusia ni</i>	6.5 to 13	0.10 to 0.20	Apply as required by scouting. Repeat applications at an interval sufficient to maintain control. ² Adjust the spray schedule (usually 5 to 10 days) according to insect pressures. Use a longer time interval when infestations are light to moderate; shorten the time interval for heavier infestations. Use lower rates when populations are at action thresholds, i.e., prior to infestations causing economic injury yet requiring treatment. Use higher rates when populations have reached economic injury levels.
	Two-spotted spider mite <i>Tetranychus urticae</i> and other <i>Tetranychus</i> sp. Broad mite <i>Polyphagotarsonemus latus</i>	6.5 to 13	0.10 to 0.20	Chilli thrips are foliar feeders and sufficient spray volume, to ensure thorough coverage, must be used for optimum control.
	Chilli Thrips <i>Scirtothrips dorsalis</i> Hood Western flower thrips <i>Frankliniella occidentalis</i> Melon thrips <i>Thrips palmi</i>	9.8 to 13	0.15 to 0.20	

¹ Apply the specified rate to the crop in an application volume of between 5 and 100 gallons of water (between 5 and 50 gallons of water for basil and chives) per acre. **DO NOT** apply more than 39 fl ozs of Pylon (0.6 lb ai) per acre, per crop growing cycle. Refer to the **Use Restrictions** and **Resistance Management** sections for specific application strategies.

² Allow at least 5 to 7 days between each application.

0 Day PHI

Conditions of Sale and Warranty

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND BASF'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF THE PRODUCT.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, EXEMPLARY, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

PC309

Pylon is a registered trademark of BASF.

© 2020 BASF Corporation
All rights reserved.

000241-00374.20200224.NVA 2020-04-132-0041
Based on: NVA 2012-04-132-0041

BASF Corporation
26 Davis Drive
Research Triangle Park, NC 27709


We create chemistry