

**ZOETIS INC.  
333 PORTAGE STREET, KALAMAZOO, MI, 49007**

Telephone: 269-359-4414

Customer Service: 888-963-8471

Website: [www.zoetis.com](http://www.zoetis.com)



Every effort has been made to ensure the accuracy of the information published. However, it remains the responsibility of the readers to familiarize themselves with the product information contained on the USA product label or package insert.

**RFD® LIQUID WORMER**

**Zoetis**

(pyrantel pamoate)

FOR PUPPIES AND DOGS

NADA #100-237, APPROVED BY FDA

Dogs usually find this dewormer very palatable and will lick the dose from the bowl willingly. If there is reluctance to accept the dose, mix in a small quantity of dog food to encourage consumption. It is recommended that dogs maintained under conditions of constant exposure to worm infestation should have a follow-up fecal exam within 2-4 weeks after first treatment. If your dog looks or acts sick, consult your veterinarian before treatment.

For maximum control and prevention of reinfection, it is recommended that puppies be treated at 2, 3, 4, 6, 8 and 10 weeks of age. Lactating bitches should be treated 2-3 weeks after whelping. Adult dogs kept in heavily contaminated quarters may be treated at monthly intervals to prevent *T. canis* reinfection.

**Easy to administer.**

**Indications for Use:** For the removal of large roundworms (*Toxocara canis* and *Toxascaris leonina*) and hookworms (*Ancylostoma caninum* and *Uncinaria stenocephala*) in dogs and puppies. The presence of these parasites should be confirmed by laboratory fecal exam. Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism.

RFD Liquid Wormer may also be used to prevent reinfection of *T. canis* in puppies and adult dogs and in lactating bitches after whelping.

**Shake Well Before Use**

**Active Ingredient:** 2.27 mg of pyrantel base as pyrantel pamoate per mL.

**Recommended Storage:** Store below 30°C (86°F).

**Warning:** Keep out of reach of children.

**Dosage and Administration:** Administer 1 teaspoon (5 mL) for each 5 lb of body weight. To assure proper dosage, weigh animal prior to treatment. It is not necessary to withhold food prior to treatment.

Distributed by: Zoetis Inc., Kalamazoo, MI 49007

Made in USA

Net Contents:	
2 fl oz (60 mL)	7985000 13909900

**CPN:** 3690185.2