

TREATS AND FEEDS 5,000 SQUARE FEET

Lawn Pro[®]

"Available only at authorized
Scotts[®] Lawn Pro[®] retailers."

Scotts

Weed and Feed
Weed Control Plus
Lawn Fertilizer



**Controls
Dandelions**
and over 50 other
broadleaf weeds

Thicker, greener turf
— satisfaction guaranteed

ACTIVE INGREDIENTS	1.18%
2,4-D [®]	0.50%
Metolachlor [®]	0.68%
OTHER INGREDIENTS	98.24%
Total	100.00%

NCAS #94757
EPA Reg. No. 530-270

EPA Est. No. 530-014

KEEP OUT OF REACH OF CHILDREN


CAUTION

See back for precautionary statements.

NET WEIGHT 14.63 lb (6.63 kg)

26-0-3

TREATS
AND FEEDS
5,000
SQUARE
FEET



Scotts[®]
Lawn Pro[®]

Weed and Feed
Weed Control Plus Lawn Fertilizer



51105

1

For best results, apply a
Scotts Annual Program
(see back for details)

Scotts.

Value-Priced Annual Program®

To improve your lawn, apply the Scotts Lawn Pro® Value-Priced Annual Program. Start anytime, just check the month and apply products at two month intervals.

* For an even better lawn... consider Scotts Best Annual Program, which provides more feeding and weed control. Your Lawn Pro® retailer can help you with this selection.



Early Spring

February - April
Before mowing and after winterizing, apply your lawn. Use the Plus Lawn Fertilizer Control Plus Lawn Fertilizer® with Weed Control for Spring Greening®. Use Scotts Lawn Pro® Value-Priced Annual Program (Scotts®) for Spring Greening®.

Late Spring

April - June
After mowing and before mowing, apply your lawn. Use the Plus Lawn Fertilizer Control Plus Lawn Fertilizer® with Weed Control for Spring Greening®. Use Scotts Lawn Pro® Value-Priced Annual Program (Scotts®) for Spring Greening®.

Summer

June - August
After mowing, apply your lawn. Use the Plus Lawn Fertilizer Control Plus Lawn Fertilizer® with Weed Control for Spring Greening®. Use Scotts Lawn Pro® Value-Priced Annual Program (Scotts®) for Spring Greening®.

Early Fall

August - November
Before mowing and after mowing, apply your lawn. Use the Plus Lawn Fertilizer Control Plus Lawn Fertilizer® with Weed Control for Spring Greening®. Use Scotts Lawn Pro® Value-Priced Annual Program (Scotts®) for Spring Greening®.

Late Fall

September - November
After mowing and before mowing, apply your lawn. Use the Plus Lawn Fertilizer Control Plus Lawn Fertilizer® with Weed Control for Spring Greening®. Use Scotts Lawn Pro® Value-Priced Annual Program (Scotts®) for Spring Greening®.



How and when to apply DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
This product applies 1.5 pounds of 2-4-D per acre per application. The maximum number of broadcast applications per treatment site is two per year.

APPLICATION

- Apply any time weeds are actively growing (when daytime temperatures are consistently above 60°F).
- Apply on a calm day.
- Apply to moist grass. To help particles stick to weed leaves the grass should be thoroughly watered or wet from dew before applying Lawn Pro Weed and Feed Weed Control Plus Lawn Fertilizer.
- Do not use this product on flowers, shrubs, fruit or vegetable plants or on creeping beangrass, St. Augustine, Ippia and dichondra lawns. On bermudagrass lawns, under certain conditions, some temporary discoloration may result.

WATERING OR RAIN

Do not water your lawn for 24 hours after application. Water will wash the weed control off the weed's leaves and reduce the effectiveness of this product. This product should not be applied if rain is expected within the next 24 hours.

IMPORTANT

WHAT TO EXPECT

Weed stems and leaves will begin to wilt and curl in a few days, but a month or more may be required for complete control. This product will not prevent weed seeds from producing new plants.

EXTREME TEMPERATURES

Avoid applying when temperatures are consistently above 90 degrees.

SEEDING/SODDING

Wait until the new grass has been mowed at least three times before applying Lawn Pro Weed and Feed Weed Control Plus Lawn Fertilizer. If you are planning to seed, you should wait at least one month after you have applied Lawn Pro Weed and Feed Weed Control Plus Lawn Fertilizer.

Advantages

- Safe feeding - won't burn when applied as directed
- Quick-greening - usually in 5 to 7 days
- Long-lasting - feeds up to 2 months
- Dependable - traditional Scotts quality
- Controls more than 50 weeds...

aster	curly dock	lambquarter	ragweed
bindweed	dandelion	lipola	scarlet
black medic	dichondra	mallow	plainspinner
bristly oxeye	dollarweed	mowbray	shadblow
blackfoot	English daisy	morningglory	shepherdspurse
bull thistle	evanscraggrass	mustard	smartweed
hackberry	filaree	penstemon	sorrel
burmudagrass	flaxweed	pepperweed	stachyris
carpetweed	ground ivy	pigweed	thymefat
centaury	heald	plains	sandwort
chicory	heald	prickly lettuce	vench
chickweed	horseweed	puncturevine	white clover
cinqfoil	knorweed	puerule	yellow rock
crabweed	(weeding)	pusley, Florida	

SPREADER

Be careful when applying this product near shrubs, flower beds or vegetable gardens in contact with weed control may damage these desirable plants. The Scotts Spreaders provide excellent controlled application to protect these areas. When using rotary spreaders, take special care near desirable plants.

Scotts No-Quibble Guarantee®

If for any reason you, the consumer, are not satisfied after using this product, you are entitled to get your money back. Simply send us evidence of purchase, and we will mail you a refund check promptly.

Dedicated To A Beautiful World®

Trusted by generations of homeowners, Scotts is dedicated to helping you enhance and protect the natural beauty of your home environment.

Scotts®

Spreading Tips

Apply header strips. Then apply product in longest direction of lawn. For best results, use a Scott's® spreader.

Spreader Settings

Scotts AccelGreen® Drop Spreaders	5 1/2
Scotts PrecisionGreen® Drop Spreaders	5
Scotts SpeedyGreen® & EdgeGuard® Rotary Spreaders	3 1/2

DIRECTIONS

Most lawns first; apply 2 header strips; apply longest direction of lawn, overlapping wheel marks.
Most lawns first; apply 2 header strips; apply longest direction of lawn, overlapping wheel marks.
Make each pass about 5 feet from previous pass to overlap spread width.

Additional Spreader Settings

EZ Drop, Republic (#530)	4
EZ Broadcast, Republic (#540)	2 1/2
Scotts Broadcast Tractor Spreader	2 1/2

(Non-Scotts spreader settings are approximate.)

KEEP OUT OF REACH OF CHILDREN CAUTION

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals: CAUTION: Harmful if swallowed. Avoid inhalation and contact with skin, eyes, or clothing. Avoid contamination of feed or foodstuffs. Do not contaminate water used for irrigation or domestic purposes.
Do not allow people (other than the applicator) or pets on treatment area during application. Do not enter treatment areas until spray has dried or dust has settled.
When using this product, wear long-sleeved shirt, long pants, socks and shoes. After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water.

FIRST AID

If in eyes: Hold eyes open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after first 5 minutes, then continue rinsing eye. **If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. **If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip water of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person. **If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance. They give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling poison control center or doctor, or going for treatment.

Environmental Hazards: Do not apply when weather conditions favor drift from target area. Do not contaminate water used for irrigation or domestic purposes. Drift or runoff may adversely affect nontarget plants. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Do not contaminate water when disposing of equipment washwaters.
SPRINKLE: Store in clean, dry place. Reveal opened bag by folding top down and securing. **GENERAL:** If Empty, Do Not reuse this container. Place in trash or other for recycling if available. **If Full:** Follow your local solid waste agency or 1-800-CLEANUP for disposal instructions. **Never** pour or sweep product into any drain or outdoor drain.

Lawn Pro® Weed and Feed Weed Control Plus Lawn Fertilizer

26-0-3

F-643

Guaranteed Analysis

Total nitrogen (%)	26%
7.7% ammoniacal nitrogen	
18.3% urea nitrogen	
0.7% other water soluble nitrogen*	
0.7% water insoluble nitrogen	
Soluble phosphorus (P ₂ O ₅)	3%
Sulfur (S)	10%
100% combined sulfur (S)	

Derived from: urea, methylbenzothio, potassium sulfate and ammonium sulfate.

*Contains 5.6% slowly available methylbenzothio and diethylthioammonium nitrogen.

Product of U.S.A.

U.S. Patent No. 5,102,440

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.regulatory-info-usa.com>

For Additional Questions, Comments and Information, contact the Scotts Lawn Pro retailer where you bought this Product or you can...




Call Scotts toll-free Customer Helpline: 1-800-545-TURF (8873)



Visit Scotts Made Easy® Web Site at www.scotts.com

Your Do-It-Yourself Lawn Care Headquarters

Scotts Lawn Pro® retailers can provide you with the do-it-yourself expertise and products to have a beautiful Scotts lawn safely and easily...for less money than hiring a lawn service. That's why we call them Lawn Pro® retailers!



©2004, Licensed by The Scotts Company. World rights reserved. Made in U.S.A.

The Scotts Company
14111 Scotsdown Rd.
Marysville, OH 43041