

Scotts.

4 STEP PROGRAM



Improves Lawn's Ability to Absorb Water & Nutrients

Covers 5,000 SQ FT

KEEP OUT OF REACH OF CHILDREN CAUTION
See back for additional precautionary statements.

ACTIVE INGREDIENT:

NOT FOR NEWLY SEEDED AREAS

28-0-7

NET WT. 13.36 lb (6.06 kg)






4STEP PROGRAM



Crabgrass Preventer plus Lawn Food



Apply in Early Spring: FEB | MAR | APR



Crabgrass Preventer plus Lawn Food Apply in Early Spring: FEB | MAR | APR







It's Easy for You to Have a Better Lawn



Start anytime

You can begin the Annual Program anytime of the year. Just check the month and select the appropriate STEP® listed. Apply the remaining steps at two-month intervals. If weeds are not a problem, apply STEP® 4 Fall Lawn Food.

Simple and easy
No more guesswork. You simply apply the right product – one STEP® at a time.

FIOTIDE APPLICATIONS
On Centipacel same, do not exceed rate of 2 lbs of Nitrogen per 1,000 sq. ft. in the north and 3 lbs of Nitrogen per 1,000 sq. ft. in the central and south. On Bahia lawns, do not exceed rate of 3 lbs of Nitrogen per 1,000 sq. ft. in the north.
South Florida is north of Ocala. Central Florida is north of Ocala to a line extending from Vero Beach to Tampa. South Florida is includes the remaining southern portion of the state.

SPREADING TIPS

Apply header strips. Then apply in the longest direction of lawn. For best results, use a Scotts® spreader. Shut off spreader or stop each time you reach a header strip. Covers 5,000 sq. ft. This package applies 2.67 lbs of product per 1,000 sq. ft.

Spreader Settings Directions

Spreader Settings

Scotts® Broadcast **Rotary Spreaders**



Drop Spreaders

Make sure spreader wheel completely overlaps previous



For additional spreader settings please call 1–800–543–TURF (8873

GET EXPERT ADVICE FOR YOUR LAWN NOW!

Want information on what to use and when? Sign up today to get FREE expert advice specifically for your lawn.

Consumer Helpline 1-800-543-TURF

Visit our website www.scotts.com



Well-fed lawns are more efficient at using available water.

Feeding with Scotts® lawn food helps grow a thick lawn with more roots to better absorb water and nutrients than an unfed lawn. This helps protect your lawn

DIRECTIONS FOR USE It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

APPLICATION:
STEP®1 Crabprass Preventer plus Lawn Food should be applied when your lawn is dry, it is not necessary to water it in after application. However, for best results the lawn should receive ¼ to ¼ inch of water from rainfall or watering within 2–3 days after application.
Apply in early spring before the 3rd or 4th mowing or before temperatures are regularly in the Apply in early spring before the 3rd or 4th mowing or before temperatures are regularly in the Apply in a construction of the performance of this product.
This product applies 1-5 lbs of pendimethalin per acre. Do not apply more than twice a year.

This product applies 1.5 lbs of pendimethalin per acre. Do not apply more than twice a year.
Not for use on dichondra or bentgrass lawns.
Do not seed or sod for 4 months following application. The Step 1 Crabgrass Preventer Plus Lawn
Food in the soil will prevent grass seed from sprouting.
Apply to prevent germination of crabgrass, foxtal, barnyardgrass, fall panicum, spurge* and oxalis
(common yellow woodsorrel):

in early spring (before dandelions reach the puffball stage)

before March in southern California

before mid-March in norther California

*For control a repeat treatment is recommended after 6–8 weeks.

Apply in August or September to prevent germination of Poa annua and chickweed. Do not enter or allow others to enter the treated area until dust has settled

Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.

Important: Do not rake or aerify for 4 months after application to avoid disturbing the layer of soil to which the SIEP® I Crobinss Preventer plus I sum food was applied.

SEDING/SODING/SPRIGGING:
Please do not apply to newly seed do a sodded a feasa until after the fourth mowing. On newly sprigged areas, wait 5 months before application.

ACTIVELY GROWING CRABGRASS:
This product controls crabgrass in its early stages of growth. This product does not control mature crabgrass and other existing weeds. You can kill broadleaf weeds with STEP® 2 Weed Control plus Lawn Food.

KEEP OUT OF REACH OF CHILDREN CAUTION PRECAUTIONARY STATEMENTS Hazards to Humans and Domestic Animals CAUTION: Hard if absorbed through the skin, Causes moderate

TRIST AID

FIRST AID

FIRST SHORT of the poper and rince slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after first 5 minutes, then continue rinsing eye.

FOR SYND OR Q.O.H.NIE as off contaminated clothing, Rince skin immediately with plenty of water for 15-20 minutes. The SYNLLOWED PLOW is a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow, Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an anaestories record.

Saming now of particles to prevent staining and unwalled tulexorp into the recovery of the prevent staining and unwalled tulexorp into the recovery of the prevent of the p

Information regarding the content and levels of metals in this product is available on the internet at http://www.regulatory-info-sc.com

Questions / Information

SMG Sold and Guaranteed by:
Listed The Scotts Company
LIST 14111 Scottslawn Rd.
Country Street Street Scotts Company LLC.
World rights reserved.

STEP® 1 Crabgrass Precenter Plus Lawn Food 28-0-7 F643

Guaranteed Analysis
Total nitrogen (N). 28%
7.2% ammoniacal nitrogen
1.7% urea nitrogen
8.3% other water soluble nitrogen*
Soluble portach (K, Soluble nitrogen*
1.0% combined sulfur (S) 7%
Soluble portach (K, Soluble nitrogen*
1.0% combined sulfur (S) 11%
11.0% combined sulfur (S

32247 39181