

# Solo-Jec<sup>®</sup> Feline 3

## Feline Rhinotracheitis-Calici-Panleukopenia Vaccine Modified Live Virus

### Description

Solo-Jec Feline 3 is for the vaccination of healthy, susceptible cats 8 weeks and older to aid in the prevention of Feline Rhinotracheitis, Calicivirus, and Panleukopenia.

### Features

Protects against:

Feline Rhinotracheitis – an upper respiratory infection spread through the air and through direct contact with other infected cats.

Calicivirus – a highly contagious infection that causes upper respiratory disease

Panleukopenia – a highly contagious disease also known as feline distemper, that may be fatal within the first five days of infection.

### Benefits

Convenient, ready-to-use package

Protection from multiple diseases

Safe for cats 8 weeks and older

### Advantages

Protection for three highly contagious feline diseases in a 1 mL dose.

### Dosage & Administration

The dosage is 1 mL injected subcutaneously using aseptic technique. Open syringe by twisting or tapping the cap against a hard surface to break the heat weld. Prepare the vaccine by injecting the diluent into the vial containing the desiccated vaccine cake. Shake well. Withdraw entire contents into the syringe.

Subcutaneous administration: Lift the loose skin behind the neck and insert needle. Inject entire contents of syringe. Do not inject directly into blood vessel. **Important note:** Before injecting vaccine, pull back slightly on syringe plunger. If blood enters the syringe freely, choose another injection site. Vaccinate kittens at 8 weeks of age or older. Administer a second 1 mL dose 3-4 weeks later. Annual revaccination is recommended.

### Precautions

Do not vaccinate pregnant animals. Only vaccinate healthy animals. Burn containers and unused contents. Although rare, severe allergic reactions (anaphylactoid) may occur that require immediate veterinary care. Antidote: Epinephrine.

### Uses

Solo-Jec Feline 3 is for subcutaneous vaccination of healthy cats eight weeks of age or older, as an aid in the prevention of disease caused by feline rhinotracheitis, calici and panleukopenia viruses.

### Product Information

Name	Code	Size
Solo-Jec <sup>®</sup> Feline 3	153-061	25x1 dose display



### Highlights

- Protection for every cat
- Everything you need for vaccination is right in the bag
- Safe for cats 8 weeks of age and older

1 mL

## Sterile Diluent

Boehringer Ingelheim Vetmedica, Inc.  
St. Joseph, MO 64506  
U.S. Vet. Lic. No. 124

Set:

80101-03

1 Dose  
Rehydrate with 1 mL

## Feline Rhinotracheitis-Calici-Panleukopenia Vaccine

Modified Live Virus

See outer package for directions and precautions.  
Store at 35-45°F (2-7°C).  
Burn this vial.

Boehringer Ingelheim Vetmedica, Inc.  
St. Joseph, MO 64506  
US Vet. Lic. No. 124

Set:  
Exp.:

153001-01



## Feline Rhinotracheitis-Calici-Panleukopenia Vaccine

Modified Live Virus

### For Use in Cats Only

Boehringer Ingelheim Vetmedica, Inc.  
St. Joseph, MO 64506  
U.S. Veterinary License No. 124

This package contains one (1 dose) vial of dry vaccine, one (1 mL) vial of vaccine diluent, and one disposable syringe.

**Indications:** For subcutaneous vaccination of healthy cats, eight weeks of age or older, as an aid in the prevention of disease caused by feline rhinotracheitis, calici and panleukopenia viruses.

**Storage:** Store in the dark at 35°-45°F (2°-7°C). Avoid freezing.

**Preservatives:** Neomycin and polymyxin B.

**SOLO-JEC® FELINE 3**

**Directions and Dosage:** The dosage is 1 mL injected subcutaneously using aseptic technique. Open syringe by twisting or tapping the cap against a hard surface to break the heat weld. Prepare the vaccine by injecting the diluent into the vial containing the desiccated vaccine cake. Shake well. Withdraw entire contents into the syringe. **Subcutaneous administration:** Lift the loose skin behind the neck and insert needle (see illustration). Inject entire contents of syringe. Do not inject directly into blood vessel. *Important note: Before injecting vaccine, pull back slightly on syringe plunger. If blood enters the syringe freely, choose another injection site.* Vaccinate kittens at 8 weeks of age or older. Administer a second 1 mL dose 3-4 weeks later. Annual revaccination is recommended.



**Precautions: Do not vaccinate pregnant animals.** Only vaccinate healthy animals. Burn containers and unused contents. Although rare, severe allergic reactions (anaphylactoid) may occur that require immediate veterinary care. Antidote: Epinephrine.



153002-00

Set:  
Exp.:

Boehringer Ingelheim Vetmedica, Inc.  
2621 North Belt Highway • St. Joseph, MO 64506-2002  
Information 800-325-9167 • Sales Service 800-325-9167 • Fax 816-326-2717

©2013 Boehringer Ingelheim Vetmedica, Inc. Solo-Jec is a registered trademark of Boehringer Ingelheim Vetmedica, Inc.

BI 1530-6 12/13