

ADM Animal Nutrition™
Safe-Guard®* Cattle and Swine
Dewormer 860200



A Supplement for Deworming Swine and Cattle

SWINE - FOR THE REMOVAL OF:

Lungworms: *Metastrongylus apri* and *Metastrongylus pudendotectus*
Gastrointestinal worms: Adult and larvae (L₃, L₄ stages, liver, lung, intestinal forms) large roundworm (*Ascaris suum*); nodular worms (*Oesophagostomum dentatum*, *O. quadrispinulatum*); small stomach worm (*Hyostromylus rubidus*); adult and larvae (L_{2, 3, 4} stages - intestinal mucosal forms) whipworm (*Trichuris suis*)
Kidneyworms: Adult and larvae *Stephanurus dentatus*.

CATTLE - FOR THE REMOVAL AND CONTROL OF:

Lungworm: (*Dictyocaulus viviparus*)
Stomach worms: Barberpole worm (*Haemonchus contortus*), Brown stomach worm (*Ostertagia ostertagi*), Small stomach worm (*trichostrongylus axei*)
Intestinal worms: Hookworm (*Bunostomum phlebotomum*), Thread-necked intestinal worm (*Nematodirus helvetianus*), Small intestinal worms (*Cooperia punctata* & *C. oncophora*)
Bankrupt worm: (*Trichostrongylus colubriformis*)
Nodular worm: (*Oesophagostomum radiatum*)

ACTIVE DRUG INGREDIENT

Fenbendazole(227 mg/lb).....0.05%

GUARANTEED ANALYSIS

Crude Protein, Min.	14.0%
Lysine, Min.	0.65%
Crude Fat, Min.	2.0%
Crude Fiber, Max.	15.0%
Calcium (Ca), Min. 0.5% Max.	1.0%
Phosphorus (P), Min.	0.25%

INGREDIENTS

Grain Products, Processed Grain By-Products, Roughage Products, Plant Protein Products, Calcium Carbonate, Sodium Propionate (A Preservative), Lignin Sulfonate.

**SEE BACK OF TAG FOR
 USE AND MIXING DIRECTIONS AND
 WARNING STATEMENT**

Fenbendazole sourced from Safe-Guard® (fenbendazole), a registered trademark of Intervet Inc., d/b/a Merck Animal Health, a subsidiary of Merck & Co. Inc., All rights reserved.

00116

**ADM Animal Nutrition, a division of
 Archer Daniels Midland Company, Quincy, IL 62305-3115
 Net Wt 50 lb (22.67 kg) or as Invoiced**

FEEDING DIRECTIONS

SWINE: Mix 200 lb of Safeguard Cattle and Swine Dewormer with 1800 lb of feed ingredients to manufacture a feed containing 22.7 mg/lb of fenbendazole. Feed this mixture as the sole ration at the rate of 18 lb per 100 pound bodyweight over a 3-12 day period to provide a total of 4.1 mg of fenbendazole per pound of bodyweight (9 mg/kg).

Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism. Swine maintained under conditions of constant worm exposure may require retreatment within four weeks after the first treatment due to reinfection.

CATTLE: For a one-day therapeutic treatment for dairy and beef cattle, mix 1 lb of Safe-Guard Dewormer 860200 per 100 lb of body weight into the daily ration. Feed as the sole ration for one day to provide 227 mg of fenbendazole per 100 lb of body weight (5 mg/kg).

Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism. No milk withdrawal required.

WARNING: Cattle must not be slaughtered within 13 days following last treatment. For dairy cattle, there is no milk withdrawal period. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

860200

00116