

SEDGE ENDER®

TRUSTED SINCE 1926

BONIDE®

Ready to Spray



Kills yellow nutsedge, crabgrass, dandelions and other listed weeds

For use on warm season and
cool season grasses

ACTIVE INGREDIENTS:

	By Wt.
Sulfentrazone: N-[2,4-dichloro-5-[4-(difluoromethyl)-4,5-dihydro-3-methyl-5-oxo-1H-1,2,4-triazol-1-yl]phenyl]methanesulfonamide..	1.36%
Proflumicarb: [N3,N3-Di-n-propyl-2,4-dinitro-6-(trifluoromethyl)-m-phenylenediamine]	2.73%

OTHER INGREDIENTS:..... 95.91%

TOTAL:..... 100.00%

Contains 0.36 pounds of active ingredient (0.12 lb ai sulfentrazone and 0.24 lb ai proflumicarb) per gallon.

EPA Est. No. 4-NY-1 EPA Reg. No. 279-3427-4

Keep Out Of Reach Of Children

CAUTION (See additional precautionary information contained inside label booklet)

Net Contents 1 Pint (473 mL)



**Weed deadly
lawn friendly**

070

SEEDGE ENDER[®] Ready to Spray

- ✓ Starts working on contact
- ✓ Rainproof - won't wash away
- ✓ Kills the weeds, not the lawn
- ✓ Kills the toughest weeds, down to the root
- ✓ Water-based

For preemergence and early postemergence control of crabgrass and selected annual grasses. Selective preemergence control of annual broadleaf weeds and preemergence and postemergence control of sedges in lawns.

EPA Est. No. 4-NY-1

EPA Reg. No. 279-3427-4

©Bonide Products, Inc. All Rights Reserved.

Distributed by:
Bonide Products, Inc.
6301 Sutliff Road
Oriskany, NY 13424
Pint

BN



DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

PRODUCT INFORMATION

Turf Use Instructions

This product is a selective preemergence and postemergence herbicide for control of crabgrass, annual grass and broadleaf weeds and sedges in residential lawns.

This product is a flowable formulation (suspension concentrate) containing 0.4 lbs. of active ingredient per gallon. The mode of action of this product involves uptake by both weed roots and shoots of existing weeds and prevention of growth by newly emerged weed seedlings.

APPLICATION INSTRUCTIONS

FOUR STEPS TO APPLY

1. **Measure** the *total* (entire) lawn area to be sprayed. For rectangular areas, multiply the length by the width to obtain *total* square feet.
2. **TO BEGIN** spraying: Connect a garden hose to the hose-end sprayer. Make sure the control valve on the nozzle is in the **OFF** position. Turn on water at faucet. Point (Aim) nozzle toward lawn and turn the control valve to the **ON** position. Water will automatically mix with the product. Spray evenly over the measured area.
3. **SPRAY:** Spray evenly over the measured area. Walk at a steady pace. This 16 fl. oz. container will treat 5,800 sq. ft. for Cool Season grasses, 2,900 sq. ft. for Warm Season Grasses and 4,000 sq. ft. for St. Augustine grass.
4. **TO STOP** spraying: Turn valve to **OFF** position. Turn off water at faucet. To relieve (discharge) pressure, turn control valve to **ON** position and wait until water stops spraying before removing sprayer from hose.

See Table 1 for specific use instructions for either cool or warm-season turfgrass varieties. For preemergence applications, spray recommended rate evenly over entire lawn. For post emergence applications, spray weeds directly until well covered. Do not over apply as this may cause undesirable turf injury.

SPECIFIC USE INSTRUCTIONS IN TURFGRASS

Established Turf

This product may be used on seeded, sodded or sprigged turfgrasses that are well established. First application of this product can be made following the second mowing providing the grass has developed into a uniform stand with a good root system. Turf injury may result from application of this product on turf that is not well established or has been weakened by stresses such as unfavorable weather conditions, disease, chemical or mechanical influences.

When applied as directed under the conditions described, the following established grasses are tolerant to this product at the recommended use rate. Do not exceed 8.25 fl. oz./1,000 sq. ft. per calendar year.

Table 1. Tolerant grasses.

Grass Type	Recommended Use Rates Do not exceed maximum use rates per turfgrass species in a single application
Cool Season Grasses	Area Covered (based on 16 fl. oz. retail container)
Fescue, fine ¹ (<i>Festuca rubra</i>), Ryegrass, perennial (<i>Lolium perenne</i>), Bluegrass, Kentucky (<i>Poa pratensis</i>) Fescue, tall ² (<i>Festuca arundinacea</i>) Cool-season mixtures	5,800 sq. ft.
Warm Season Grasses	
Bahia grass (<i>Paspalum notatum</i>) Buffalograss (<i>Buchloe dactyloides</i>) Carpetgrass (<i>Axonopus affinis</i>) Centipedegrass (<i>Eremochloa ophiuroides</i>) Kikuyugrass (<i>Pennisetum clandestinum</i>) Seashore Paspalum (<i>Paspalum vaginatum</i>) Zoysiagrass ³ (<i>Zoysia japonica</i>) Bermudagrass ⁴ (<i>Cynodon dactylon</i>) & hybrids	2,900 sq. ft.

Grass Type	Recommended Use Rates Do not exceed maximum use rates per turfgrass species in a single application
Warm Season Grasses	Area Covered (based on 16 fl. oz. retail container)
St. Augustine grass	4,000 sq. ft.

¹ Use of this product on certain varieties of Chewing Fine Fescue or Tall Fescue cultivars may result in undesirable injury.

² This product may be applied more than once per year if needed as long as total amount applied does not exceed the maximum application rate listed per turf type or 8.25 fl. oz./1,000 sq. ft. per calendar year, whichever is lower.

³ This product application may cause temporary discoloration to exposed leaf surfaces on certain cultivars of zoysiagrass. Treated turfgrass will recover with new growth. Discolored leaf tissue will be removed with mowing. To reduce potential for discoloration, do not apply this product on turfgrass that is weakened by weather, mechanical, chemical, disease or other related stress. Maintain proper cultural practices such as adequate moisture and fertility levels to promote healthy turf growth.

⁴ Sequential application of this product may be needed in bermudagrass lawns for season-long control of yellow nutsedge. If sequential application is needed apply no more than 16 fl. oz. to 5,800 sq. ft.

This product has demonstrated tolerance on both cool and warm season turfgrasses. However, not all varieties and or cultivars have been evaluated. Turfgrass managers desiring to treat newly released varieties should first apply this product to a small area prior to treatment of larger areas.

WEED CONTROLLED WITH PREEMERGENCE APPLICATION

Table 2.

Common Name	Scientific Name
Barnyardgrass	<i>(Echinochloa crus-galli)</i>
Bluegrass annual ¹	<i>(Poa annua)</i>
Carpetweed	<i>(Mollugo verticillata)</i>
Carolina Geranium	<i>Geranium carolinianum</i>
Chickweed, common ²	<i>(Stellaria media)</i>
Chickweed, mouseear (from seed)	<i>(Cerastium vulgatum)</i>
Crabgrass (Large, Smooth) ³	<i>(Digitaria spp.)</i>
Crowfootgrass	<i>(Dactyloctenium aegyptium)</i>
Cupgrass, Woolly	<i>(Eriochloa villosa)</i>
Foxtails, Annual	<i>(Setaria spp.)</i>
Goosegrass ⁴	<i>(Eleusine indica)</i>
Hairy Bittercress	<i>Cardamine hirsute</i>
Henbit ²	<i>(Lamium amplexicaule)</i>
Itchgrass	<i>(Rottboellia exaltata)</i>
Knotweed, prostrate	<i>(Polygonum aviculare)</i>
Kochia	<i>(Kochia scoparia)</i>
Kyllinga, green	<i>(Kyllinga brevifolia)</i>
Kyllinga, false	<i>(Kyllinga gracillima)</i>
Lambsquarters, common	<i>(Chenopodium album)</i>
Lovegrass	<i>(Eragrostis spp.)</i>
Nutsedge, yellow	<i>(Cyperus esculentus)</i>
Panicum (Texas, Fall)	<i>(Panicum spp.)</i>
Pigweed, Redroot	<i>(Amaranthus retroflexus)</i>
Purslane, Common	<i>(Portulaca oleracea)</i>
Pusley, Florida	<i>(Richardia scabra L.)</i>
Sedge, globe	<i>(Cyperus globulosus)</i>

WEED CONTROLLED WITH PREEMERGENCE APPLICATION CONT.

Common Name	Scientific Name
Sedge, cylindrical	<i>(Cyperus retrorsus)</i>
Sedge, Surinam	<i>(Cyperus surinamensis)</i>
Sedge, Texas	<i>(Cyperus polystachyos)</i>
Shepherdspurse ²	<i>(Capsella bursa-pastoris)</i>
Signalgrass, broadleaf	<i>(Brachiaria playphylla)</i>
Speedwell	<i>(Veronica spp.)</i>
Sprangletop	<i>(Leptochola spp.)</i>
Spurge, prostrate	<i>(Euphorbia humistrata)</i>
Witchgrass	<i>(Panicum capillare L.)</i>
Woodsorrel, yellow (from Seed)	<i>(Oxalis stricta)</i>

1. Poa annua is a winter annual. Applications of this product should be made in August or September to established, non overseeded turf before Poa Annua germinates. Consult local Cooperative Extension Service for more specific applications timings for your local area since August and September are approximate timings.

2. Apply this product in late summer, fall or winter before weed germinates for control.

3. Crabgrass control in cool season grasses from fall application: Areas where the ground is subject to winter freezes, this product may be applied in the fall at rates listed in Table 1 (5,800 sq. ft./16 fl. oz.) for cool season grasses when soil temperatures are below 50°F, but prior to the ground freezing. This will provide control of crabgrass the following spring.

4. In areas of extended growing season and heavy pressure weed control will be enhanced by second application of this product 60-90 days after initial application.

Postemergence Control of Annual, Biennial & Perennial Broadleaf Weeds, Grasses and Sedges

This product will control or suppress the weeds listed in Table 3 when applied shortly after weeds have emerged. Do not exceed the application rate specified for the turf species in Table 1. Do not apply adjuvants or surfactants with post applications of this product unless tested on small areas prior to broadcast applications.

WEEDS CONTROLLED OR SUPPRESSED WHEN APPLICATION MADE POSTEMERGENCE TO WEEDS

Table 3.

Common Name	Scientific Name	Controlled	Suppressed
Bittercress	(<i>Cardamine spp.</i>)		X
Black medic	(<i>Medicago lupulina</i>)		X
Buttercups	(<i>Ranunculus spp.</i>)		X
Carolina geranium	(<i>Geranium carolinianum</i>)	1	
Carpetweed	(<i>Mollugo verticillata</i>)		X
Chickweed, common	(<i>Stellaria media</i>)	1	
Chickweed, mouseear	(<i>Cerastium vulgatum</i>)		X
Cinquefoil	(<i>Potentilla spp.</i>)		X
Clover	(<i>Trifolium spp.</i>)		X
Crabgrass (Large and Smooth)	(<i>Digitaria spp.</i>)	2	
Cudweed	(<i>Gnaphalium spp.</i>)		X
Dandelion	(<i>Taraxacum officinale</i>)		X
Dock, Curly	(<i>Rumex crispus</i>)		X
Evening primrose	(<i>Oenothera biennis</i>)		X
Fiddleneck	(<i>Amsinckia spp.</i>)		X
Filaree	(<i>Erodium spp.</i>)		X
Goldenrod	(<i>Solidago spp.</i>)	1	
Goosegrass	(<i>Eleusine indica</i>)	2	
Ground ivy	(<i>Glechema hederacea</i>)	1	
Henbit	(<i>Lamium amplexicaule</i>)		X
Knotweed, prostrate	(<i>Polygonum aviculare</i>)		X
Kochia	(<i>Kochia scoparia</i>)	1	
Lambsquarters, common	(<i>Chenopodium album</i>)		X
Lawn burweed	(<i>Soliva pterosperma</i>)		X
Lespedeza, common	(<i>Lespedeza striata</i>)		X
Mallow, common	(<i>Malva neglecta</i>)		X

WEEDS CONTROLLED OR SUPPRESSED WHEN APPLICATION MADE POSTEMERGENCE TO WEEDS CONT.

Common Name	Scientific Name	Controlled	Suppressed
Nutsedge, Yellow	<i>(Cyperus esculentus)</i>	1	
Parsley piert	<i>(Alchemilla arvensis)</i>	1	
Pigweed, Redroot	<i>(Amaranthus retroflexus)</i>	1	
Pigweed, Tumble	<i>(Amaranthus albus)</i>	1	
Pineapple weed	<i>(Matricaria matricarioides)</i>		X
Plantain, buckhorn	<i>(Plantago lanceolata)</i>	1	
Puncture weed	<i>(Tribulus terrestris)</i>		X
Purslane, common	<i>(Portulaca oleracea)</i>		X
Pusley, Florida	<i>(Richardia scabra)</i>	1	
Redweed	<i>(Melochia corchorifolia)</i>		X
Rocket, London	<i>(Sisymbrium irio)</i>		X
Smartweed, Pennsylvania	<i>(Polygonum pensylvanicum)</i>	1	
Sorrel, Red	<i>(Rumex acetosella)</i>		X
Speedwell	<i>(Veronica spp.)</i>	1	
Spurge, (annuals)	<i>(Euphorbia spp.)</i>	1	
Spurge, prostrate	<i>(Euphorbia humistrata)</i>	1	
Spurge, spotted	<i>(Euphorbia maculata)</i>	1	
Star of Bethlehem	<i>(Ornithogalum umbellatum)</i>	1	
Velvetleaf	<i>(Abutilon theophrasti)</i>		X
Violet, wild	<i>(Viola pratincola)</i>		X
Wild garlic	<i>(Allium vineale)</i>		X
Wild onion	<i>(Allium canadense)</i>		X
Woodsorrel, creeping	<i>(Oxalis corniculata)</i>		X
Woodsorrel, yellow	<i>(Oxalis stricta)</i>	1	

1. Weeds are suppressed at lower label rates (5,800 sq. ft./16 fl. oz.). For optimum control apply rates of at 2,900 sq. ft. coverage/16 fl. oz. in a single application. Do not exceed the application rate specified for the turf species in Table 1.
2. This product controls crabgrass and goosegrass when applied postemergent to newly emerged weeds in the 1-4 leaf stage of development.

Application to Reseeded, Overseeded or Sprigged Areas

Reseeding, overseeding or sprigging of treated areas within three to six (3-6) months after application of this product could inhibit the establishment of desirable grasses. See table for exact intervals from application.

Best results are obtained for reseeding or overseeding when mechanical or power seeding equipment (slit seeders) are used to give good seed to soil contact and proper soil cultivation, irrigation and fertilization practices are followed.

OVERSEEDING INTERVALS AFTER PRODUCT APPLICATIONS

Table 4.

Total Amount of This Product Applied Square feet per 16 fl. oz.	Interval (Months Before Overseeding)		
	North	Transition	South
5,800 sq. ft.	3	3	3
4,000 sq. ft.	3	3	3
2,900 sq. ft.	4	4	4

Other Use Precautions

- Do not use on turfgrasses other than those listed on this label.
- Do not graze or feed livestock, forage cut from areas treated with this product.
- Do not apply directly to landscape ornamentals or ornamental beds.
- Do not apply to areas where dichondra, colonial bentgrass, velvet bentgrass or annual bluegrass (*Poa annua*) are desirable species.
- Do not apply to newly set sod until the sod has rooted and exposed edges have grown in.
- Do not apply with adjuvants or surfactants.

STORAGE AND DISPOSAL

Pesticide Storage: Store and transport in upright position. Store only in original container, in a cool dry place inaccessible to children or pets.

Pesticide Disposal: *If partly filled* – Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

Container Disposal: *If empty* – Non-refillable container. Do not reuse or refill this container. Offer for recycling, if possible.

FIRST AID

IF INHALED	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.• Call a <u>poison control center or doctor for further treatment advice.</u>
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a <u>poison control center or doctor for treatment advice.</u>
IF IN EYES	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a <u>poison control center or doctor for treatment advice.</u>
IF SWALLOWED	<ul style="list-style-type: none">• Call a <u>poison control center or doctor immediately for treatment advice.</u>• Do not give any liquid to the person.• Do not induce vomiting unless told to do so by the poison control center or doctor.• Do not give anything by mouth to an unconscious person.

HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact the National Pesticide Information Center at 1-800-858-7378 for emergency medical treatment information.

PRECAUTIONARY STATEMENTS

Hazards to Humans (and Domestic Animals)

CAUTION: Causes moderate eye irritation. Harmful if inhaled, swallowed, or absorbed through skin. Avoid breathing vapor or spray mist. Avoid contact with skin, eyes or clothing. Prolonged or frequently repeated skin contact while mixing or handling may cause allergic reactions in certain individuals. Wear long-sleeved shirt and long pants, socks, shoes, and chemical-resistant gloves (such as natural rubber, selection Category A). Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

Environmental Hazards

This product is toxic to marine/estuarine invertebrates. Drift and runoff may be hazardous to terrestrial and aquatic plants in neighboring areas. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

Groundwater advisory: Sulfentrazone is known to leach through soil into groundwater under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

Do not use on coarse soils classified as sand which have less than 1% organic matter.

WARRANTY

Seller warrants that this product complies with the specifications expressed in this label. To the extent consistent with applicable law, Seller makes no other warranties, and disclaims all other warranties, express or implied, including but not limited to warranties of merchantability and fitness for the intended purpose. To the extent consistent with applicable law, Seller's liability or default, breach or failure under this label shall be limited to the amount of the purchase price. To the extent consistent with applicable law, Seller shall have no liability for consequential damages.

For information on pesticide products (including health concerns, medical emergencies, or pesticide incidents) call National Pesticide Information Center at 1-800-858-7378.