

Specimen Label



Snapshot[®] DG

SPECIALTY HERBICIDE

®™ Trademarks of Dow AgroSciences, DuPont or Pioneer and their affiliated companies or respective owners

A selective preemergence herbicide for control of certain broadleaf weeds and annual grasses in:

- Landscape Ornamentals
- Christmas Tree Plantations
- Container Grown Ornamentals
- Field Grown Ornamentals
- Groundcovers/Perennials
- Non-Bearing Fruit and Nut Trees
- Non-Bearing Vineyards
- Non-Cropland

Active Ingredients:

trifluralin: α, α, α -trifluoro-2,6-dinitro- <i>N</i> , <i>N</i> -dipropyl- <i>p</i> -toluidine.....	2.0%
isoxaben: <i>N</i> -[3-(1-ethyl-1-methylpropyl)-5- isoxazolyl]-2,6-dimethoxybenzamide and isomers.....	0.5%
Other Ingredients.....	97.5%
Total.....	100.0%

Contains 1.25 lb active ingredient per 50 lb bag.

Precautionary Statements

Hazards to Humans and Domestic Animals

EPA Reg. No. 62719-175

Keep Out of Reach of Children

CAUTION

Causes Moderate Eye Irritation • Harmful If Swallowed Or Inhaled • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reaction In Some Individuals

Avoid breathing dust or spray mist and contact with eyes or clothing. Wash thoroughly with soap and water after handling.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.

User Safety Recommendations (Cont.)

- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Users should remove PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a Poison Control Center or doctor for treatment advice.

If swallowed: Call a Poison Control Center or doctor immediately for treatment advice. Have a person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a Poison Control Center or doctor. Do not give anything by mouth to an unconscious person.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a Poison Control Center or doctor for further treatment advice.

Have the product container or label with you when calling a Poison Control Center or doctor or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

Environmental Hazards

This pesticide is extremely toxic to freshwater marine, and estuarine fish and aquatic invertebrates including shrimp and oyster. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply in a manner which will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements of this label about personal protective equipment (PPE), and restricted entry interval. The requirements in the box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Waterproof gloves
- Shoes plus socks

Non-Agricultural Use Requirements

The requirements of this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

For applications in landscape settings and in non-cropland sites, do not enter or allow others to enter the treated area until dusts have settled.

Storage and Disposal

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store in original container only. In case of spill, contain material and dispose as waste.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Nonrefillable rigid containers 5 gallons or less:

Container Handling: Nonrefillable container. Do not reuse or refill this container.

Storage and Disposal (Cont.)

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Nonrefillable nonrigid containers:

Container Handling: Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available, or dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Refillable rigid containers larger than 5 gal:

Container Handling: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Nonrefillable rigid containers larger than 5 gal:

Container Handling: Nonrefillable container. Do not reuse or refill this container.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

General Information

Snapshot® DG specialty herbicide is a preemergence product for control of certain broadleaf weeds and annual grasses in container and landscape ornamentals, nursery stock, Christmas trees, groundcovers/perennials, non-bearing fruit and nut trees, non-bearing vineyards and non-cropland areas. Apply Snapshot DG anytime prior to germination of target weeds, or immediately after cultivation. Length of weed control will vary with rate of Snapshot DG applied, weed population, potting media or soil conditions, temperature, watering regime, and other factors. Following application, user should monitor and observe level of weed control over time to determine when additional applications may be needed.

General Use Precautions and Restrictions

In Arizona: The state of Arizona has not approved Snapshot DG for use on plants grown for agricultural/commercial production; such as on Christmas tree plantations, container grown or field grown ornamentals, non-bearing fruit and nut trees and non-bearing vineyards.

Snapshot DG controls weeds growing from seed. Snapshot DG does not control established weeds, weeds growing from stolons, rhizomes, or root pieces. Existing weeds should be controlled by cultivation or with postemergence herbicides. Weed residues, prunings and trash should be removed or thoroughly mixed into soil prior to treatment. Soil should be in good condition and free of clods at the time of application. A single rainfall or sprinkler irrigation of 0.5 inches or more, or flood irrigation is required to activate Snapshot DG. Optimum weed control is obtained when Snapshot DG is activated within 3 days of application. If rainfall or irrigation does not occur within 3 days of application and tillage is possible, Snapshot DG may be activated using cultivation equipment capable of uniformly mixing the herbicide into the upper 1 to 2 inches of soil. Failure to activate Snapshot DG within 3 days of application may result in erratic control of annual grasses. Do not apply when wind conditions favor drift of Snapshot DG granules from the target area.

Repeat applications at 150 lb per acre and higher should not be made sooner than 60 days after a previous application of Snapshot DG. Do not apply over 600 pounds per acre total of Snapshot DG within a 12-month period.

Tolerance of Turfgrass Adjacent to Ornamental Plantings: Accidental application to turf may occur when applying Snapshot DG to ornamental plantings. Snapshot DG is not recommended for weed control in turfgrasses, but turfgrasses are generally tolerant to small amounts of this product that fall outside of the intended area of application.

Do not aerially apply Snapshot DG.

Treatment Species Not Listed on the Label for Snapshot DG

Users who wish to use Snapshot DG on plant species not recommended on this label may determine the suitability for use by treating a small number of such plants at a recommended rate. Prior to treatment of larger areas, the treated plants should be observed for any sign of herbicidal injury for during 30 to 60 days of normal growing conditions to determine if the treatment is safe to the target plant species. The user assumes responsibility for any plant damage or other liability resulting from use of Snapshot DG on plant species not recommended on this label.

Application Instructions

Apply Snapshot DG using a drop or rotary-type spreader designed to apply granular herbicides or insecticides. Calibrate application equipment prior to use according to manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly. Do not use spreaders that apply material in narrow concentrated bands. Avoid skips or overlaps as poor weed control or crop injury may occur. More uniform application may be achieved by spreading half of the required amount of product over the area and then applying the remaining half in swaths at right angles to the first.

Application Techniques for Applying Snapshot DG

- When using a drop-type spreader, a splashboard mounted under the hopper will provide more even granule distribution.
- A chain fastened to the side of the spreader and allowed to drag on the soil surface can be used to mark the edge of the treated swath and help prevent skips or overlaps.
- For treating smaller areas or rows of nursery stock or ornamental beds, a hand held or push-type rotary applicator such as a whirlybird or cyclone unit is recommended. For hand held units, walk and turn the crank at a constant rate of speed.
- A shaker-type applicator made from a small container with holes punched in the bottom is recommended for small, difficult to treat areas. Carefully measure the amount of product needed to avoid over application.
- The weight-to-volume conversion table below provides equivalent amounts of Snapshot DG.

Broadcast Rates

Rate lb/Acre	Amount/1000 sq ft		Amount/100 sq ft	
	(lb)	(quarts)	(lb)	(cups)
100	2.30	1.75	0.23	0.7
150	3.45	2.60	0.35	1.0
200	4.60	3.50	0.46	1.4

1 lb of Snapshot DG = 0.75 quart = 3 cups

Spreader Settings as a Guide for Calibration

Note: These settings are provided as a suggested starting point in calibrating each individual spreader. Since no two spreaders are alike, these settings are not intended to be used as absolute recommendations by Dow AgroSciences or the spreader manufacturer.

Settings for Warren T-7 II Spreader			
Target Rate of Snapshot DG			
Speed (mph)	100 lb/acre	150 lb/acre	200 lb/acre
2.0	2 1/2	3 1/4	3 3/4
2.5	3	3 1/2	4 1/4
3.0	3 1/4	4	5

Warren spreader settings are displayed in half-number increments only. The suggested settings that indicate quarter settings can be accomplished by placing the calibration arm between the half number marks on the spreader.

Weeds Controlled or Suppressed

Weeds controlled when applied at 100 lb per acre (2.3 lb per 1000 sq ft):

Common Name
 aster, slender
 barnyardgrass
 bluegrass, annual
 bursage, annual
 celery, wild
 chickweed, common
 clover, white
 crabgrass
 cudweed, purple
 cupgrass, southwestern
 fiddleneck, coast
 filaree, redstem
 fleabane, blackleaved
 fleabane, dwarf
 foxtail, yellow
 groundcherry, lanceleaf
 henbit
 horseweed
 junglerice
 knotweed, prostrate
 lambsquarters, common
 mallow, little
 mustard, Indian
 mustard, wild
 nightshade, black
 oat, wild
 panicum, fall
 pepperweed, Virginia
 pigweed
 pineappleweed
 plantain, slender
 purslane, common
 radish, wild
 ragweed, common
 rocket, London
 rockpurslane, desert
 shepherdspurse
 sibara
 smartweed, Pennsylvania
 sowthistle, annual
 speedwell, purslane
 sprangletop, Mexican
 telegraphplant
 thistle, Russian
 witchgrass

Scientific Name
Aster exilis
Echinochloa crus-galli
Poa annua
Ambrosia acanthicarpa
Apium leptophyllum
Stellaria media
Trifolium repens
Digitaria spp.
Gnaphalium purpureum
Eriochloa gracilis
Amsinckia intermedia
Erodium cicutarium
Conyza bonariensis
Conyza ramosissima
Setaria glauca
Physalis lanceifolia
Lamium amplexicaule
Conyza canadensis
Echinochloa colonum
Polygonum aviculare
Chenopodium album
Malva parviflora
Brassica juncea
Sinapis arvensis
Solanum nigrum
Avena fatua
Panicum dichotomiflorum
Lepidium virginicum
Amaranthus spp.
Matricaria matricarioides
Plantago elongata
Portulaca oleracea
Raphanus raphanistrum
Ambrosia artemisiifolia
Sisymbrium irio
Calandrinia ciliata
Capsella bursa-pastoris
Sibara virginica
Polygonum pennsylvanicum
Sonchus oleraceus
Veronica peregrina
Leptochloa univervia
Heterotheca grandiflora
Salsola iberica
Panicum capillare

Weeds controlled when applied at 150 lb per acre (3.5 lb per 1000 sq ft):

Common Name
 aster, heath
 bittercress
 bittercress, hairy
 brassbuttons, southern
 carrot, wild
 chamberbitter
 chickweed, mouseear
 dandelion
 eclipta
 fireweed
 foxtail, giant
 galinsoga, hairy
 geranium, Carolina
 groundsel, common
 ladysthumb
 lettuce, prickly
 lovegrass
 mallow, dwarf
 marestalk
 mayweed
 morningglory, ivyleaf
 mustard, black
 pennywort

Scientific Name
Aster ericoides
Cardamine oligosperma
Cardamine hirsuta
Cotula australis
Daucus carota
Phyllanthus urinaria
Cerastium vulgatum
Taraxacum officinale
Eclipta prostrata
Erechtites hieracifolia
Setaria faberi
Galinsoga ciliata
Geranium carolinianum
Senecio vulgaris
Polygonum persicaria
Lactuca serriola
Eragrostis spp.
Malva rotundifolia
Hippuris vulgaris
Anthemis cotula
Ipomoea hederacea
Brassica nigra
Hydrocotyle spp.

Common Name (Cont.)

phyllanthus, long-stalk
 plantain, bracted
 plantain, broadleaf
 plantain, buckhorn
 pokeweed, common
 rockpurslane, redmaids
 ryegrass, annual
 sida, prickly
 sorrell, red
 speedwell, thymeleaf
 spurge, hyssop
 spurge, spotted
 sweetclover, yellow
 tansymustard, green
 velvetleaf
 woodsorrel, yellow

Scientific Name

Phyllanthus tenellus
Plantago aristata
Plantago major
Plantago lanceolata
Phytolacca americana
Calandrinia ciliata var. *menziesii*
Lolium multiflorum
Sida spinosa
Rumex acetosella
Veronica serpyllifolia
Euphorbia hyssopifolia
Euphorbia maculata
Melilotus officinalis
Descurainia pinnata
Abutilon theophrasti
Oxalis stricta

Weeds controlled when applied at 200 lb per acre (4.6 lb per 1000 sq ft):

Common Name
 barley, hare
 bromegrass
 burclover, California
 cheat
 datura
 dogfennel
 eveningprimrose
 fescue, rattail
 filaree, whitestem
 goosefoot, nettleleaf
 goosegrass
 jimsonweed
 knotweed, silversheath
 Kochia
 medic, black
 mullein, turkey
 nettle, burning
 nettle, stinging
 oxtongue, bristly
 pimpernel, scarlet
 sandbur, field
 signalgrass
 sowthistle, spiny
 spurge, petty
 spurge, prostrate
 stinkgrass
 sunflower
 swinecress
 thistle, musk
 willoweed, panicle
 woodsorrel, creeping

Scientific Name

Hordeum leporinum
Bromus spp.
Medicago polymorpha
Bromus secalinus
Datura spp.
Eupatorium capillifolium
Oenothera spp.
Vulpia myuros
Erodium moschatum
Chenopodium murale
Eleusine indica
Datura stramonium
Polygonum argyrocoleon
Kochia scoparia
Medicago lupulina
Eremocarpus setigerus
Urtica urens
Picris dioica
Picris echinoides
Anagallis arvensis
Cenchrus incertus
Brachiaria spp.
Sonchus asper
Euphorbia peplus
Euphorbia humistrata
Eragrostis cilianensis
Helianthus spp.
Coronopus didymus
Carduus nutans
Epilobium paniculatum
Oxalis corniculata

Weeds partially controlled or suppressed when applied at 200 lb per acre (4.6 lb per 1000 sq ft):

Common Name
 bindweed, field
 carpetweed
 dock, curly
 johnsongrass (seedling)
 mallow, Venice
 milkweed, honeyvine
 morningglory, tall
 panicum, Texas
 pusley, Florida
 shattercane

Scientific Name

Convolvulus arvensis
Mollugo verticillata
Rumex crispus
Sorghum halepense
Hibiscus trionum
Ampelamus albidus
Ipomoea purpurea
Panicum texanum
Richardia scabra
Sorghum bicolor

Uses

Ornamental Plantings

Snapshot DG is recommended as a preemergence treatment for control of certain broadleaf weeds and annual grasses in container grown ornamentals, landscape ornamentals, field grown ornamentals, ground covers/perennials, Christmas trees, non-bearing fruit and nut crops, and non-bearing vineyards. Apply Snapshot DG anytime prior to germination of target weeds, or immediately after cultivation.

Refer to the General Information section of this label for General Use Precautions and information on application, application rates and weeds controlled.

Special Use Precautions:

To avoid possible injury, do not apply Snapshot DG to:

- nursery, forest, or Christmas tree: seedling beds, cutting beds, or transplant beds.
- nursery seedbeds or forest or Christmas tree seedling transplant beds.
- unrooted liners or cuttings that have been planted in pots for the first time.
- pots less than four inches wide.
- bedding plants or areas where bedding plants will be planted or transplanted within one year following application.
- groundcovers until they are established and well rooted.

Do not apply Snapshot DG to newly transplanted ornamentals, nursery stock, Christmas trees, groundcovers, non-bearing fruit and nut trees and non-bearing vineyards until soil or potting media has been settled by packing and irrigation or rainfall and no cracks are present or injury may occur.

Repeat applications at 150 lb per acre and higher should not be made sooner than 60 days after a previous application of Snapshot DG. Do not apply over 600 lb per acre total of Snapshot DG within a 12-month period.

Where loss of stand has occurred in field grown ornamentals, use untreated soil as fill around roots when replacing plants or injury may occur.

Note: Injury on the following plant species has been observed following applications of Snapshot DG and use is not recommended.

Common Name	Scientific Name
ballerina cranesbill	<i>Geranium cinereum</i> 'Ballerina'
bertram anderson lingwort	<i>Pulmonaria longifolia</i>
bugle	<i>Ajuga</i> spp.
cajeput tree	<i>Melaleuca quinquenervia</i>
candytuft	<i>Iberis</i> spp.
Carolina rhododendron	<i>Rhododendron carolinianum</i>
dwarf burning bush	<i>Euonymus alatus</i> 'compacta'
foxglove	<i>Digitalis purpurea</i>
green yucca	<i>Yucca recurvifolia</i>
hydrangea	<i>Hydrangea</i> spp.
luxuriant bleeding heart	<i>Dicentra luxuriant</i>
mountain sandwort	<i>Arenaria montana</i>
mustard	<i>Brassica</i> spp.
oak leaf acanthus	<i>Acanthus mollis</i>
prince of wales juniper	<i>Juniperus horizontalis</i> 'Prince of Wales'
purple coneflower	<i>Echinacea purpurea</i>
roseum elegans	<i>Rhododendron catawbiense</i> 'Roseum elegans'
rhododendron	
spurge	<i>Euphorbia</i> spp.
stonecrop	<i>Sedum</i> spp.
summer phlox	<i>Phlox paniculata</i>
white festival baby's breath	<i>Gypsophila paniculata</i>
wine periwinkle	<i>Vinca minor</i> 'Atropurpurea'

Snapshot DG may be used on the following established plant species (Note: Limitations on recommended treatment methods):

Trees	Recommended Treatment Method: C = Container Grown F = Field Grown	
Scientific Name	Common Name	
<i>Abies balsamea</i>	fir, balsam	C, F
<i>Abies concolor</i>	fir, white	F
<i>Abutilon hybridum</i>	albus-flowering maple	C, F
	luteus-flowering maple	C, F
	roseus-flowering maple	C, F
	tangerine-flowering maple	C, F
	vesuvius red-flowering maple	F
<i>Acer ginnala</i>	flame maple	F
<i>Acer rubrum</i>	red maple	F
	red sunset maple	F
<i>Acer saccharinum</i>	silver maple	C, F
<i>Alsophila australis</i>	Australian tree fern	C, F
<i>Areacastrium romanzoffianum</i>	queen palm	C, F
<i>Betula nigra</i>	birch, river	C, F
<i>Betula papyrifera</i>	paper birch	F
<i>Brachychiton populneus</i>	bottle tree	C, F
<i>Bucida buceras</i>	black olive	F
<i>Ceratonia siliqua</i>	carob	F
<i>Cercis canadensis</i>	redbud	C, F
<i>Chamaecyparis obtusa</i>	filicoides-fernspray cypress	F
<i>Chamaecyparis obtusa</i>	gracilis-slender hinoki cypress	F
<i>Chamaecyparis pisifera</i>	sawara-false cypress	F
	squarrosa-moss cypress	F
<i>Chamaedorea cataractarum</i>	cat palm	F
	palm	C, F
<i>Chamaedorea costaricana</i>	palm	C, F
<i>Chamaedorea elegans</i>	parlor palm	C, F
<i>Cornus florida</i>	cloud nine dogwood	C, F
	dogwood, flowering	C, F
<i>Crataegus viridis</i>	green hawthorn	F
<i>Cryptomeria japonica</i>	cryptomeria, Japanese	C, F
<i>Cupaniopsis anacardioides</i>	carrot wood	F
<i>Cupressocyparis x 'Emerald Isle'</i>	emerald island leyland cypress	C, F
<i>Cupressus arizonica</i>	Arizona cypress	C, F
<i>Cupressus glabra</i>	Arizona cypress	F
<i>Cupressus sempervirens</i>	Italian cypress	C, F
<i>Dicksonia antarctica</i>	Tasmanian tree fern	C, F
<i>Elaeagnus angustifolia</i>	Russian olive	C, F
<i>Elaeagnus x 'Gilt edge'</i>	gilt edge silverberry	C, F

Trees (Cont.)

**Recommended Treatment Method:
C = Container Grown
F = Field Grown**

Scientific Name	Common Name	
<i>Eucalyptus camaldulensis</i>	red gum eucalyptus	F
<i>Eucalyptus cinerea</i>	eucalyptus, mealy	F
	silver dollar eucalyptus	F
<i>Eucalyptus microtheca</i>	coolibah tree	C, F
<i>Eucalyptus sideroxylon</i>	eucalyptus, red ironbark	F
<i>Ficus benjamina</i>	fig	C, F
	mini ficus	C, F
<i>Fraxinus udhei</i>	shamel ash	C, F
<i>Ginkgo biloba</i>	ginkgo (maidenhair tree)	F
<i>Gleditsia triacanthos</i>	honey locust	F
	shademaker honey locust	F
<i>Heteromeles arbutiflora</i>	toyon	F
<i>Illicium floridanum</i>	Florida anise-tree	C, F
<i>Juniperus virginiana</i>	redcedar, eastern	C, F
<i>Liquidambar styraciflua</i>	sweetgum, American	F
<i>Magnolia grandiflora</i>	magnolia, southern	C, F
<i>Morus alba</i>	white mulberry	F
<i>Musa aluminata</i>	banana	C, F
<i>Oxydendrum arboreum</i>	sourwood	C, F
<i>Picea abies</i>	pendula-weeping Norway spruce	C, F
	repens-spreading Norway spruce	C, F
	spruce, Norway	C, F
<i>Picea glauca</i>	conica-dwarf Alberta spruce	C, F
<i>Picea glauca conica</i>	dwarf Alberta spruce	F
<i>Picea pungens</i>	dwarf globe blue spruce	C, F
	glauca-Colorado blue spruce	C, F
	hoopsii-hoop's blue spruce	C, F
	koster-koster blue spruce	F
<i>Pinus aristata</i>	bristlecone pine	F
<i>Pinus canariensis</i>	canary island pine	F
<i>Pinus contorta</i>	shore pine, beach pine	F
<i>Pinus eldarica</i>	eldarica pine	C, F
<i>Pinus leucodermis</i>	Bosnian pine	C, F
<i>Pinus mugo</i>	pumilio-shrubby swiss	C, F
	mountain pine	
<i>Pinus nigra</i>	Austrian black pine	C, F
<i>Pinus radiata</i>	monterey pine	F
<i>Pinus strobus</i>	eastern white pine	C, F
<i>Pinus strobus</i>	white pine	C, F
<i>Pinus sylvestris</i>	columnar Scotch pine	C, F
	Scotch pine	C, F
<i>Pinus thunbergiana</i>	Japanese black pine	C, F
<i>Platanus occidentalis</i>	American sycamore	F
<i>Platanus racemosa</i>	Californina sycamore	F
<i>Podocarpus</i> spp.	podocarpus	F
<i>Populus deltoides</i>	cottonwood	F
<i>Prosopis chilensis</i>	Chilean mesquite	C, F
<i>Prunus yedoensis</i>	yoshino flowering cherry	F
<i>Quercus ilicifolia</i>	bear oak	F
<i>Quercus palustris</i>	pin oak	F
<i>Quercus phellos</i>	willow oak	C, F
<i>Quercus rubra</i>	red oak	C, F
<i>Quercus virginiana</i>	live oak	C, F
<i>Salix babylonica</i>	babylon weeping willow	F
	corkscrew willow	F
<i>Sequoiadendron giganteum</i>	giant sequoia	F
<i>Swietenia mahogani</i>	mahogany	F
<i>Tabebuia caraiiba</i>	yellow tab	F
<i>Tsuga canadensis</i>	eastern hemlock	C, F
<i>Ulmus parvifolia</i>	Chinese elm	F
<i>Washingtonia robusta</i>	Mexican fan palm	F

Shrubs

**Recommended Treatment Method:
C = Container Grown
F = Field Grown**

Scientific Name	Common Name	
<i>Abelia grandiflora</i>	edward goucher abelia	C, F
	glossy abelia	C, F
<i>Acacia abyssinica</i>	abyssinica acacia	C, F
<i>Acacia redolens</i>	acacia, prostrate	C, F
<i>Acacia stenophylla</i>	shoestring acacia	C, F
<i>Acalypha wilkesiana</i>	copper leaf	C, F
<i>Acer ginnala</i>	amur maple	C, F
<i>Acer palmatum</i>	coral bark Japanese maple	C, F
	dwarf Japanese maple	C, F
<i>Agapanthus africanus</i>	queen anne lily of the Nile	C, F
<i>Agave americana</i>	century plant	F
<i>Anisodonteia hypomandarum</i>	cape mallow	C, F
<i>Aptenia cordifolia</i>	red apple aptenia	C, F
<i>Ardisia japonica</i>	chirimen marlberry	C, F
<i>Astible arendsii</i>	false spiraea	C, F
<i>Astilbe chinensis</i>	astilbe/false spirea	C, F

Shrubs (Cont.)

Recommended Treatment Method
C = Container Grown
F = Field Grown

Scientific Name	Common Name	
<i>Athyrium nipponicum</i>	Japanese painted fern	C, F
<i>Baccharis pilularis</i>	coyotebush	F
<i>Berberis gladywomensii</i>	william penn barberry	C, F
<i>Berberis mentorensis</i>	mentor barberry	C, F
<i>Berberis thunbergii</i>	aurea-golden Japanese barberry	F
	crimson pygmy barberry	C, F
	rose glow barberry	C, F
	atropurea-redleaf Japanese barberry	F
	cherry bomb barberry	C, F
<i>Bougainvillea</i> spp.	barbara karst	C, F
	California gold	C, F
	pink pixie	C, F
	scarlet o'hara	C, F
	temple fire	C, F
	Texas dawn	C, F
<i>Buxus</i> x 'Green velvet'	green velvet boxwood	C, F
<i>Buxus microphylla japonica</i>	boxwood, Japanese	C, F
<i>Buxus microphylla Koreana</i>	Korean boxwood	F
<i>Buxus sempervirens</i>	boxwood, common	C, F
<i>Callistemon citrinus</i>	bottlebrush, lemon	F
<i>Callistemon viminalis</i>	weeping bottlebrush	C, F
<i>Calluna vulgaris</i>	spring torch Scotch heather	C, F
<i>Camellia japonica</i>	camellia	C, F
<i>Caryopteris x clandonen</i>	dark knight bluebeard	C, F
<i>Cassia artemisioides</i>	cassia, feathery	C, F
<i>Ceanothus</i> spp.	wild lilac	F
<i>Cephalotaxus drupacea</i>	plum yew	C, F
<i>Cerastium tomentosum</i>	snow-in-summer	C, F
<i>Chamaecyparis obtusa</i>	kosteri cypress	C, F
	nana-dwarf hinoki cypress	C, F
	torulosa cypress	C, F
<i>Chamaecyparis pisifera</i>	baileyi-dogwood	F
	flaviramea-dogwood	F
	squarrosa minima cypress	C, F
<i>Chamaecyparis pisifera</i> spp.	filifera-thread cypress	C, F
<i>Chrysalidocarpus lutescens</i>	areca palm	F
<i>Clethra alnifolia</i>	summersweet	C, F
<i>Cleyera japonica</i>	cleysera, Japanese	C, F
<i>Coleonema pulchrum</i>	pink breath of heaven	C, F
<i>Convolvulus cneorum</i>	bush morning glory	C, F
<i>Cornus alba</i>	sibirica-Siberian dogwood	C, F
<i>Cornus stolonifera</i>	baileyi-red-osier dogwood	F
	flaviramea-yellowtwig dogwood	F
<i>Cotinus coggygria</i>	royal purple smoke tree	C, F
<i>Cotinus dammeri</i>	coral beauty smoke tree	C, F
	eichholz smoke tree	C, F
<i>Cotoneaster adpressus</i>	praecox-early cotoneaster	C, F
<i>Cotoneaster apiculatus</i>	cotoneaster, cranberry	C, F
<i>Cotoneaster congestus</i>	cotoneaster, Pyrenees	C, F
<i>Cotoneaster dammeri</i>	cotoneaster, bearberry	C, F
<i>Cotoneaster himalayana</i>	Himalayan cotoneaster	C, F
<i>Cotoneaster horizontalis</i>	cotoneaster, rock	C, F
<i>Cotoneaster opiculata</i>	cotoneaster	C, F
<i>Cycas revoluta</i>	sago palm	C, F
<i>Cytisus praecox</i>	hollandia-warminster broom	C, F
<i>Cytisus scoparius</i>	lena-Scotch broom	C, F
<i>Cytisus</i> spp.	holandia-Scotch broom	F
<i>Daphne odora</i>	fragrant daphne	C, F
<i>Deutzia crenata</i>	nakiana-dwarf deutzia	C, F
<i>Deutzia gracilis</i>	slender gracilis	C, F
<i>Dodonea viscosa</i>	hopseed bush	F
<i>Elaeagnus pungens</i>	fruitland silver berry	C, F
<i>Erica cinerea</i>	purple bell heather	C, F
<i>Erica vagans</i>	cornish heather	C, F
<i>Erica x darleyensa</i>	Mediterranean pink heather	C, F
<i>Eugenia myrtifolia</i>	dwarf brush cherry	C, F
<i>Euonymus</i> x 'Aureo variegatus'	gold spot euonymus	C, F
<i>Euonymus</i> x 'Chollipo'	chollipo euonymus	C, F
<i>Euonymus fortunei</i>	canadale gold euonymus	C, F
	emerald'n gold euonymus	F
	sunspot euonymus	C, F
<i>Euonymus japonica</i>	silver king euonymus	F
	variegated evergreen euonymus	C, F
<i>Euonymus kiatschovica</i>	spreading euonymus	C, F
<i>Euonymus vegetus</i>	bigleaf wintercreeper	C, F
<i>Euryops pectinatus</i>	dwarf euryops	C, F
<i>Fatsyhedera japonica</i>		C, F
<i>Fatsia japonica</i>	Japanese aralia	C, F
<i>Felicia ameloides</i>	blue marguerite	C, F
<i>Forsythia intermedia</i>	forsythia, border	C, F
<i>Forsythia</i> x 'Spring glory'	spring glory forsythia	C, F
<i>Gardenia jasminoides</i>	august beauty gardenia	C, F

Shrubs (Cont.)

Recommended Treatment Method
C = Container Grown
F = Field Grown

Scientific Name	Common Name	
	gardenia	C, F
	radican gardenia	C, F
<i>Gaultheria shallon</i>	salal/lemon leaf	C, F
<i>Gelsemium sempervirens</i>	Carolina jessamine	C, F
<i>Genista pilosa</i>	woadwaxen	C, F
<i>Hibiscus rosa-sinensis</i>	ross estey-hibiscus	C, F
<i>Hibiscus syriacus</i>	rose of sharon, red bird	C, F
	rose of sharon, red heart	F
	rose of sharon, woodbridge	C, F
	rose of sharon, aphrodite	
	rose of sharon, helene	
<i>Ilex aquifolium</i>	Balkans holly	F
	gold coast holly	F
<i>Ilex aquipernyi</i>	san jose holly	C, F
<i>Ilex attenuata</i>	savannah holly	C, F
<i>Ilex cornuta</i>	burford holly	C, F
	dwarf burford holly	C, F
	needle point holly	C, F
<i>Ilex crenata</i>	compacta-dwarf Japanese holly	C, F
<i>Ilex crenata</i>	convexa holly	C, F
	dwarf Chinese holly	C, F
	green luster holly	C, F
	helleri-heller's Japanese holly	C, F
	hetzii's Japanese holly	C, F
	stokesii Japanese holly	C, F
<i>Ilex glabra</i>	compacta-compact inkberry holly	C, F
	nordica-inkberry holly	C, F
<i>Ilex meserveae</i>	blue boy holly	C, F
	blue girl holly	C, F
	China boy holly	
	China girl holly	
	ebony magic holly	F
<i>Ilex vomitoria</i>	nana-dwarf yaupon holly	C, F
	pendula-weeping yaupon holly	C, F
	yaupon holly	C, F
<i>Illicium annisatum</i>	mystery gardenia	C, F
<i>Itea ilicifolia</i>	henry garnet holly leaf sweetspire	C, F
<i>Ixora collinea</i>	ixora	C, F
<i>Juniperus chinensis</i>	hollywood juniper	C, F
	media-old gold juniper	C, F
	pfitzer juniper	C, F
	pfitzerana glauca-blue juniper	C, F
	pfitzerana-pfitzer juniper	C, F
	sea green juniper	F
	torulosa-hollywood juniper	C, F
<i>Juniperus conferta</i>	emerald sea shore juniper	C, F
	shore juniper	C, F
<i>Juniperus horizontalis</i>	andorra juniper	C, F
	bar harbor juniper	C, F
	blue chip juniper	C, F
	blue rug juniper	C, F
	creeping juniper	C, F
	dwarf andorra juniper	C, F
	huntington blue juniper	C, F
	plumosa-andorra juniper	C, F
	wiltonii-blue carpet juniper	C, F
<i>Juniperus procumbens</i>	nana-dwarf Japaneses garden juniper	C, F
<i>Juniperus prostrata</i>	prostrata juniper	C, F
<i>Juniperus sabina</i>	broadmoor juniper	C, F
	foemina-hicks juniper	C, F
	savin juniper	C, F
	tamariscifolia-tam juniper	C, F
<i>Juniperus scopulorum</i>	emerald green juniper	F
<i>Juniperus squamata</i>	blue juniper	C, F
	blue star juniper	C, F
	parsonii juniper	C, F
<i>Kalmia latifolia</i>	laurel, mountain	C, F
<i>Lagerstroemia indica</i>	crepe myrtle	C, F
<i>Lantana</i> spp.	lantana	C, F
<i>Lavandula angustifolia</i>	English lavender	C, F
<i>Lavandula latifolia</i>	English spike lavender	C, F
<i>Lavandula officianalis</i>	English lavender	C, F
<i>Leptospermum scoparium</i>	New Zealand tea tree	C, F
<i>Leucothoe axillaris</i>	leucothoe, coast	C, F
<i>Leucothoe fontanesiana</i>	leucothoe, drooping	C, F
<i>Ligustrum japonicum</i>	privet, Japanese	C, F
	wax ligustrum	C, F
	yellow tip ligustrum	C, F
<i>Ligustrum lucidum</i>	privet, glossy	C, F
<i>Ligustrum ovalifolium</i>	California privet	F
<i>Ligustrum texanum</i>	howardi privet	C, F
	wax leaf privet	C, F
<i>Ligustrum vicaryi</i>	privet, golden	F
	vicary golden privet	F

Shrubs (Cont.)

Recommended Treatment Method
C = Container Grown
F = Field Grown

Scientific Name	Common Name	
<i>Ligustrum vulgare</i>	iodense privet	C, F
<i>Livistona chinensis</i>	Chinese fountain palm	F
<i>Lonicera fragrantissima</i>	winter honeysuckle	C, F
<i>Lonicera periclymenum</i>	flowering woodbine	C, F
	serotina woodbine	C, F
<i>Lonicera sempervirens</i>	trumpet honeysuckle	C, F
<i>Loropetalum chinense</i>	fringe flower	C, F
<i>Mahonia aquifolium</i>	dwarf Oregon grape	C, F
	compacta	
<i>Mahonia bealei</i>	leather leaf mahonia	C, F
<i>Mahonia repens</i>	creeping mahonia	C, F
<i>Myrica cerifera</i>	wax myrtle	C, F
<i>Nandina domestica</i>	compacta-dwarf heavenly bamboo	C, F
	harbour dwarf-heavenly bamboo	C, F
	heavenly bamboo (nandina)	C, F
	nana compacta-heavenly bamboo	C, F
	nana purpurea-heavenly bamboo	C, F
	woods dwarf-heavenly bamboo	C, F
<i>Nerium oleander</i>	hardy red oleander	C, F
	oleander	C, F
	ruby lace oleander	C, F
<i>Osmanthus fortunei</i>	fortunes osmanthus	C, F
<i>Pachysandra terminalis</i>	Japanese spurge	C, F
<i>Phoenix roeoloenii</i>	pigmy date palm	C, F
<i>Photinia fraseri</i>	fraser's photinia	C, F
<i>Pieris japonica</i>	lily-of-the-valley	C, F
	mountain fire lily-of-the-valley	C, F
	snowdrift lily-of-the-valley	C, F
	temple bells lily-of-the-valley	C, F
	valley rose lily-of-the-valley	C, F
	valley valentine lily-of-the-valley	C, F
<i>Pieris japonica x forestii</i>	forest flame lily-of-the-valley	C, F
<i>Pinus mugo</i>	mugo-mugho pine	C, F
<i>Pittosporum tobira</i>	green pittosporum	C, F
	wheeler's dwarf pittosporum	C, F
<i>Plumbago ariculata</i>	blue cape plumbago	F
<i>Plumbago capensis</i>	plumbago	C, F
<i>Podocarpus macrophyllum</i>	yewpine	C, F
<i>Polygala dalmaisiana</i>	sweet pea shrub	C, F
<i>Polystichum</i>	tassel fern	C, F
	polyblepharum	
<i>Potentilla fragiformis</i>	cinquefoil	F
<i>Potentilla fruticosa</i>	cinquefoil	C, F
	gold drop potentilla	F
	goldfinger potentilla	C, F
	red ace potentilla	C, F
	sunset potentilla	C, F
	tangerine potentilla	C, F
	spring cinquefoil	C, F
<i>Potentilla verna</i>	dwarf pink flowering almond	C, F
<i>Prunus glandulosa</i>	lolendei monrovia pyracantha	C, F
<i>Pyracantha fortuneana</i>	monon pyracantha	F
	red elf hybrid pyracantha	F
	rutgers hybrid pyracantha	C, F
	santa cruz pyracantha	C, F
	victory pyracantha	F
<i>Raphiolepis indica</i>	charisma-monruce raphiolepis	C, F
	enchantress-moness raphiolepis	F
	raphiolepis (India hawthorn)	C, F
	springtime-monme raphiolepis	F
<i>Raphiolepis ovata</i>	roundleaf raphiolepis	C, F
<i>Rhododendron</i>	cannon's double azalea	C, F
	calendulaceum	
	flame azalea	F
	golden flare azalea	C, F
	klondike azalea	C, F
<i>Rhododendron</i>	butterfly rhododendron	F
<i>campylocarpum</i>		
<i>Rhododendron</i>	PJM rhododendron	C, F
	carolinianum x daurium	
<i>Rhododendron</i>	catawba album rhododendron	C, F
	catawba rhododendron	C, F
	lord roberts rhododendron	C, F
	rocket rhododendron	C, F
<i>Rhododendron caucasicum</i>	cunningham white rhododendron	C, F
<i>x ponticum</i>		
<i>Rhododendron exbury</i>	cannon's double azalea	C, F
	golden flare azalea	C, F
	klondike azalea	C, F
<i>Rhododendron forrestii</i>	gomer waterer rhododendron	C, F
	repens	
<i>Rhododendron forrestii x griersonianum</i>	elizabeth rhododendron	C, F

Shrubs (Cont.)

Recommended Treatment Method
C = Container Grown
F = Field Grown

Scientific Name	Common Name	
<i>Rhododendron</i>	jean marie rhododendron	C, F
	griffithianum	
<i>Rhododendron hybrid spp.</i>	America rhododendron	C, F
	English roseum rhododendron	F
	nova zembra rhododendron	C, F
	scintillation rhododendron	C, F
<i>Rhododendron impeditum</i>	rhododendron	C, F
<i>Rhododendron indica</i>	formosa azalea	C, F
<i>Rhododendron indica</i>	waucabusa azalea	C, F
<i>Rhododendron kaempferi</i>	blue danube azalea	C, F
<i>Rhododendron kerume</i>	coral bells azalea	C, F
	hino crimson azalea	C, F
	hino pink azalea	C, F
	mildred azalea	C, F
	snow azalea	C, F
<i>Rhododendron maximum</i>	rhodie max (rosebay)	C, F
<i>Rhododendron</i>	rhododendron	F
	mucronulatum	
<i>Rhododendron obtusum</i>	hino-crimson azalea	C, F
<i>Rhododendron ponticum</i>	chioniodes rhododendron	C, F
<i>Rhododendron ponticum</i>	daphnioides rhododendron	C, F
<i>Rhododendron x</i>	purple gem	
<i>Rhododendron</i>	dwarf scarlet wonder rhododendron	C, F
	racemosum	
	tribly rhododendron	C, F
	unique rhododendron	C, F
	vulcan rhododendron	C, F
<i>Rhododendron</i>	ramapo rhododendron	C, F
	sassthigiatim x	
	carolinianum	
<i>Rhododendron satuski</i>	gumpo pink azalea	C, F
	higasa azalea	F
	reijn azalea	C, F
<i>Rhododendron spp.</i>	carror azalea	C, F
	hybrids	
	fashion azalea	C, F
	gerard christina azalea	F
	girard roberta azalea	C, F
	golden flare exbury azalea	F
	helmut vogel azalea	F
	hershey red azalea	F
	hot shot azalea	C, F
	hume azalea	F
	inga azalea	F
	irene koster azalea	C, F
	president clay azalea	C, F
	tradition azalea	C, F
	sumac, African	C, F
<i>Rhus lancea</i>	ramanan rose	C, F
<i>Rosa rugosa</i>	rosemary	F
<i>Rosmarinus officinalis</i>	dusty-miller/silver ragweed	C, F
<i>Senecio cineraria</i>	Japanese skimmia	C, F
<i>Skimmia japonica</i>	reeve's skimmia	C, F
<i>Skimmia revesiana</i>	Paraguay nightshade	C, F
<i>Solanum rantonetii</i>	anthony waterer spiraea	C, F
<i>Spiraea bumalda</i>	first snow spiraea	C, F
<i>Spiraea x cinerea</i>		
	'Grefsheim'	
<i>Spiraea japonica</i>	dolchia spiraea	C, F
	Japanese alpine spiraea	C, F
	shirobana spiraea	C, F
	bridal wreath	C, F
<i>Spiraea vanhouttei</i>	Chinese lilac	C, F
<i>Syringa rothomagensis</i>	lilac, common	F
<i>Syringa vulgaris</i>	yew, Japanese	F
<i>Taxus cuspidata</i>	cape honeysuckle	C, F
<i>Tecomaria capensis</i>	ternstroemia, Japanese	C, F
<i>Ternstroemia gymnanthera</i>	emerald arborvitae	C, F
<i>Thuja occidentalis</i>	globosa-globe arborvitae	C, F
	little giant-dwarf arborvitae	C, F
	nigra-dark American arborvitae	C, F
	pyramidalis arborvitae	C, F
	rheingold arborvitae	C, F
	techny arborvitae	F
<i>Thuja occidentalis</i>	woodwardii arborvitae	C, F
<i>Thuja orientalis</i>	aureus nana-dwarf golden arborvitae	C, F
	minima glauca-dwarf arborvitae	C, F
<i>Veitchia merrilli</i>	Christmas palm	F
<i>Viburnum bodnantense</i>	pink dawn viburnum	C, F
<i>Viburnum carlesii</i>	Koreanspice viburnum	C, F
<i>Viburnum davidii</i>	david viburnum	C, F
<i>Viburnum japonicum</i>	viburnum	F
<i>Viburnum judd (v. X juddii)</i>	viburnum	C, F
<i>Viburnum lantana</i>	wayfaring tree	F

Shrubs (Cont.)

Recommended Treatment Method
C = Container Grown
F = Field Grown

Scientific Name	Common Name	
<i>Viburnum opulus sterile</i>	common snowball viburnum	F
<i>Viburnum plicatum tomentosum</i>	doublefile viburnum	C, F
<i>Viburnum setigerum</i>	tea viburnum	F
<i>Viburnum tinus compactum</i>	spring bouquet viburnum	F
<i>Viburnum trilobum</i>	cranberry bush	C, F
<i>Viburnum trilobum compactum</i>	dwarf cranberry bush	C, F
<i>Viburnum x pragense</i>	viburnum	C, F
<i>Weigela florida</i>	bristol ruby weigela	C, F
	java red weigela	C, F
	minuet weigela	C, F
<i>Xylosma congestum</i>	xylosma	F
<i>Yucca filamentosa</i>	yucca	C, F

Groundcovers/ Perennials

Recommended Treatment Method:
C = Container Grown
F = Field Grown

Scientific Name	Common Name	
<i>Achillea millefolium</i>	common yarrow	C, F
<i>Agapanthus africanus</i>	lily of the Nile	C, F
<i>Agapanthus "Peter Pan"</i>		C, F
<i>Aistroemeria aurea</i>	Peruvian lily	C, F
<i>Ammophila breviligulata</i>	beechgrass	C, F
<i>Antirrhinum majus</i>	snapdragon	C, F
<i>Arctotheca calendula</i>	cape weed	F
<i>Argyranthemum frutescens</i>	Paris daisy	C, F
<i>Artemisia schmidtiana</i>	angels' hair	C, F
<i>Asparagus retrofractus</i>	fern	C, F
<i>Asteriscus maritimus</i>	gold coin daisy	C, F
<i>Astilbe Deutschland</i>	deutschland astilbe	C, F
<i>Asparagus retrofractus</i>		C, F
<i>Asparagus variegata</i>	tree fern	C, F
<i>Aster novae-angliae</i>	New England aster	C, F
<i>Aster novi-belgii</i>	New York aster	C, F
<i>Begonia cordifolia</i>	heartleaf begonia	C, F
<i>Begonia semperflorens</i>	white ambassador begonia	C, F
<i>Bidens ferulifolia</i>	peter's gold bidens	C, F
<i>Brachycome x 'New amethyst'</i>	swan river daisy new amethyst	C, F
<i>Callistephus chinensis</i>	China aster	C, F
<i>Carex spp.</i>	sedge	C, F
<i>Carpobrotus edulis</i>	ice plant, largeleaf	F
<i>Catharanthus roseus</i>	Madagascar periwinkle	C, F
<i>Cerastium tomentosum</i>	snow in the summer	C, F
<i>Ceratostigma plumbaginoides</i>	dwarf plumbago	C, F
<i>Chrysanthemum morifolium</i>	florist's chrysanthemum	C, F
<i>Chrysanthemum sp.</i>	chrysanthemum species	C, F
<i>Clematis integrifolia caerulea</i>	blue bell clematis	C, F
<i>Clivia miniata</i>	kafir lily	C, F
<i>Coreopsis verticillata</i>	coreopsis, threadleaf	C, F
<i>Cortaderia selloana</i>	pampas grass	C, F
<i>Cuphea hyssopifolia</i>	false or Mexican heather	C, F
<i>Cyperus albostratus</i>	dwarf umbrella grass	C, F
<i>Dahlia x 'Royal dahlia pink'</i>	dwarf dahlia wendy pink	C, F
<i>Delosperma alba</i>	white iceplant	F
<i>Descampsia caespitosa</i>	descampsia	C, F
<i>Dianthus gratianopolitanus</i>	crimson treasure cheddar pink	C, F
<i>Dietes vegeta</i>	fortnight lily	C, F
<i>Drosantheum floribundum</i>	trailing rosea iceplant	F
<i>Drosantheum hispidum</i>	iceplant	C, F
<i>Ensete ventricosum</i>	red abyssinian banana	C, F
<i>Equisetum scirpoides</i>	dwarf horsetail	C, F
<i>Erianthus ravennae</i>	hardy pampasgrass	C, F
<i>Erysimum "Bowles mauve"</i>	wallflower	C, F
<i>Euryops pectinatus</i>	dwarf euryops	C, F
<i>Eustoma grandiflorum</i>	pink lisianthus	C, F
<i>Festuca ovina glauca</i>	blue fescue	C, F
<i>Fuchsia x 'Santa Claus'</i>	santa claus fuchsia	C, F
<i>Gaillardia aristata</i>	blanket flower	C, F
<i>Gaillardia grandiflora</i>	goblin blanket flower	C, F
<i>Gaura lindheimeri</i>	gaura	C, F
<i>Gazania rigens leucolaena</i>	gazania, trailing	C, F
<i>Gazania spp.</i>	gazania	C, F
<i>Geranium incanum</i>	cranesbill	C, F
<i>Geranium subcaulescens</i>	blackeyed magenta cranesbill	C, F
<i>Hakonechloa macroaureola</i>	golden hakonechloa	C, F

Groundcovers/ Perennials (Cont.)

Recommended Treatment Method:
C = Container Grown
F = Field Grown

Scientific Name	Common Name	
<i>Hedera canariensis</i>	ivy, Algerian	F
<i>Hedera helix</i>	ivy, English	C, F
<i>Helichrysum petiolatum</i>	white licorice plant	C, F
<i>Hemerocallis spp.</i>	daylily	C, F
<i>Hesperaloe parviflora</i>	red yucca	C, F
<i>Heuchera americana</i>	palace purple	C, F
<i>Heuchera micrantha</i>	coral bells	C, F
<i>Hippeastrum hybrid</i>	amaryllis	C, F
<i>Hosta 'Francee'</i>	francee plantain lily	C, F
<i>Hosta lancifolia</i>	albo-marginata hosta	C, F
<i>Hosta 'Patriot'</i>	patriot plantain lily	C, F
<i>Hymenoxys acaulis</i>	angelita daisy	C, F
<i>Hypericum spp.</i>	St. Johnswort	C, F
<i>Impatiens wallerana</i>	busy lizzie	C, F
<i>Iris pumila</i>	yellow dwarf bearded iris	C, F
<i>Iris siberica</i>	blue siberian iris	C, F
<i>Jasminum nitidum</i>	angelwing jasmine	C, F
<i>Lampranthus spectabilis</i>	trailing iceplant	F
<i>Leptospermum scoparium</i>	broom tree/tea/manuka	C, F
<i>Liatris spicata</i>	gay feather	C, F
<i>Limonium perezii</i>	statice	C, F
<i>Liriope gigantea</i>	white lily turf	C, F
<i>Liriope muscari</i>	lilac beauty lily turf	C, F
	majestic lily turf	C, F
	monroe white lily turf	C, F
	silvery sunproof lily turf	C, F
	variegated liriope lily turf	C, F
	big blue lily turf	C, F
<i>Liriope spicata</i>	green/creeping lily turf	C, F
<i>Lobelia erinus</i>	lobelia	C, F
<i>Lobularia maritima</i>	sweet alyssum	C, F
<i>Lonicera japonica</i>	honeysuckle, Japanese	F
<i>Lysimachia punctata</i>	dotted loosestrife	C, F
<i>Mathiola incana</i>	stock	C, F
<i>Miscanthus sinensis</i>	eulalia grass	C, F
<i>Monarda didyma</i>	bee balm	C, F
<i>Moraea iridioides</i>	African iris	C, F
<i>Oenothera speciosa</i>	siskiyou evening primrose	C, F
<i>Ophiopogon japonicus</i>	dwarf mondo grass	C, F
	mondo grass	C, F
<i>Osteospermum fruticosum</i>	freeway daisy	C, F
<i>Pachysandra terminalis</i>	Japanese spurge	C, F
<i>Parthenocissus quinquefolia</i>	Virginia creeper	C, F
<i>Pelargonium x hortorum</i>	zonal geranium	C, F
<i>Pelargonium peltatum</i>	ivy geranium	C, F
<i>Pennisetum alopecuroides</i>	fountain grass	C, F
<i>Pennisetum setaceum</i>	chrimson fountaingrass	C, F
<i>Pentas lanceolata</i>	star cluster	C, F
<i>Penstemon x 'Apple blossom'</i>	apple blossom penstemon	C, F
<i>Penstemon gentianoides</i>	hartwig penstemon	C, F
<i>Perovskia atriplicifolia</i>	Russian sage	C, F
<i>Petunia-hybrids</i>	garden petunias	C, F
<i>Phalaris arundinacea picta</i>	ribbon grass	C, F
<i>Ratibida columnifera</i>	Mexican hat	C, F
<i>Rudbeckia fulgida</i>	blackeyed susan	C, F
<i>Rudbeckia hirta</i>	blackeyed susan	C, F
<i>Ruellia brittoniana</i>	dwarf katie ruellia	C, F
<i>Salvia grahamii</i>	graham's sage	C, F
<i>Salvia leucantha</i>	Mexican bush sage	C, F
<i>Sedum x 'Autumn joy'</i>	autumn joy stonecrop	C, F
<i>Sedum x 'Vera jameson'</i>	vera jameson stonecrop	C, F
<i>Targetes patula 'Little hero'</i>	little hero marigold	C, F
<i>Trachelospermum asiaticum</i>	asian jasmine	C, F
<i>Tulbaghia violacea</i>	society garlic	C, F
<i>Verbena peruviana</i>	st. paul verbena	C, F
<i>Vinca major</i>	periwinkle, bigleaf	C, F
<i>Vinca minor</i>	periwinkle, dwarf	F
<i>Vinca spp.</i>	periwinkle	F
<i>Zinnia elegans</i>	dwarf zinnia	C, F

Non-Bearing Fruit and Nut Trees

Recommended Treatment Method:
C = Container Grown
F = Field Grown

Common Name	
almond	F
apple	F
apricot	F
avocado	F
blackberry	F

**Non-Bearing Fruit and Nut Trees
Non-bearing Vineyards¹
(Cont.)**

**Recommended Treatment Method:
C = Container Grown
F = Field Grown**

Common Name

blueberry	F
boysenberry	F
cherry, sour	F
cherry, sweet	F
currant	F
dewberry	F
elderberry	F
fig	F
filbert	F
gooseberry	F
grape, American	F
grape, European	F
grapefruit	F
kiwi	F
lemon	F
loganberry	F
macadamia nut	F
nectarine	F
olive	F
orange	F
peach	F
pear	F
pecan	F
pistachio	F
plum	F
pomegranate	F
prune	F
raspberry	F
walnut, black	F
walnut, English	F

¹Non-bearing fruit and nut trees and non-bearing vineyards are defined as plants which will not bear fruit for at least one year after treatment.

Ornamental Bulbs

Do not use Snapshot DG in bulb production fields. For use in landscape settings only.

Snapshot DG may be applied for control of susceptible annual weeds in ornamental bulbs, e.g., bulbous iris, daffodil (narcissus), hyacinth and tulip. Apply Snapshot DG to the soil surface 2 to 4 weeks after planting, but prior to the emergence of annual weeds. Snapshot DG may also be applied following bulb emergence but prior to flowering and after flowering. For fall planted bulbs, apply Snapshot DG again in late winter or early spring to weed-free soil surfaces.

Special Use Precautions:

Do not apply to tulip plants that have emerged to a height greater than 3/4 inch.

Do not apply to gladiolus prior to emergence or if corms are less than one inch in diameter.

Do not apply to bulbs while they are flowering.

Shadehouse Areas

Snapshot DG may be applied in open shadehouse-type structures where the natural flow of air is unimpeded. Do not apply in enclosed greenhouses or in enclosed shadehouse-type structures. Do not apply within three weeks prior to enclosing greenhouse or poly-type structures.

Non-Cropland

Snapshot DG is recommended as a preemergence treatment for control of certain broadleaf weeds and annual grasses on non-cropland areas such as industrial sites, utility substations, highway guardrails, sign posts, delineators, etc. Apply Snapshot DG anytime prior to germination of

target weeds. Areas to be treated should be free of established weeds or existing weeds should be controlled with postemergence herbicides.

Refer to the General Information section of this label for General Use Precautions and information on Application, Application Rates, and Weeds Controlled.

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

Warranty Disclaimer

Dow AgroSciences warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. Dow AgroSciences MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Dow AgroSciences or the seller. All such risks shall be assumed by buyer.

Limitation of Remedies

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Dow AgroSciences' election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

Dow AgroSciences shall not be liable for losses or damages resulting from handling or use of this product unless Dow AgroSciences is promptly notified of such loss or damage in writing. In no case shall Dow AgroSciences be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use, and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of the Seller or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

®™ Trademarks of Dow AgroSciences, DuPont or Pioneer and their affiliated companies or respective owners

**Produced for
Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis, IN 46268**

Label Code: CD02-895-020

Replaced Label: D02-895-001

EPA accepted 03/20/02