

Spectro® 90 WDG fungicide is a combination of two proven broad-spectrum actives. Both have contact and systemic properties to provide preventive and curative control to many devastating plant diseases. Spectro controls more than 65 diseases on more than 110 ornamental species, including annual and perennial flowers, bedding plants, groundcovers, trees and shrubs.

COMPETITIVE ADVANTAGES

EASY-TO-USE BROAD-SPECTRUM CONTROL

- Convenient Pre-Mixed Formula: Combines contact and systemic fungicides for effortless application.
- Rapid Action: Provides immediate protection by halting germinating spores and stopping mycelium growth on contact.
- Proven Control: Offers persistent systemic curative effects against diseases.
- Resistance Management: Acts as a rotational partner in managing resistance with two modes of action.
- Comprehensive Disease Control: Effective against a wide range of ornamental diseases.
- Tank-Mix Compatible: Easily mixes with most commonly used pesticides for added convenience.
- Chemigation: application through irrigation systems.

AT-A-GLANCE INFO

EPA REG. NO.	1001-72
ACTIVE INGR.	chlorothalonil (72%), thiophanate-methyl (18%)
FORMULATION	water dispersible granules
CHEM. FAMILY	chloronitrile, benzimidazole
FRAC NO.	1 and M5
REI	12 hours
PACKAGE SIZE	5 lb (4 per case)
APPLICATION	spray, drench

- FIELD-GROWN NURSERY
- CONTAINER-GROWN NURSERY
- **GREENHOUSE**
- DANGER



Spectro 90 WDG

TWO ACTIVE INGREDIENTS FOR POWERFUL CONTROL.

KEY USES

More than 110 ornamental species including: Annual and perennial flowers **Bedding plants** Foliage plants Groundcovers Deciduous and evergreen trees and shrubs

KEY DISEASES CONTROLLED

(See product label for complete list) Actinopelte leaf spot Alternaria leaf spot, leaf blight Anthracnose Ascochyta blight Bipolaris (Helminthosporium) leaf spot Black spot of rose Botrytis flower spot, flower blight Botrytis leaf spot, leaf blight Cephalosporium leaf spot Cercospora leaf spot Cercosporidium leaf spot Colletotrichum leaf blotch,

leaf spot, blights Corynespora leaf spot Coryneum blight, shothole

Curvularia flower spot

Cylindrosporium leaf spot

Dactylaria leaf spot

Didymellina leaf spot

Drechslera leaf spot, ink spot

Exobasidium leaf blister

Entomosporium (Fabraea) leaf spot

Fusarium leaf spot

Gloeosporium black leaf spot

Marssonina leaf spot

Monilinia blossom blight, twig blight

Mycosphaerella ray blight

Myrothecium leaf spot, brown rot

Nematostoma leaf blight

Ovulinia flower blight

Phyllsoticta leaf spot

Phytophthora aerial blight

Ramularia leaf spot

Rhizoctonia web blight

Rhizopus blossom blight

Sclerotinia flower blight

Septoria leaf spot

Sphaeropsis leaf spot

Stagonospora leaf scorch

Taphrina leaf blister

Volutella leaf blight

USE SUMMARY

Apply Spectro® 90 WDG at a rate of 1.0 to 2.0 pounds per 100 gallons of water unless other directions are given. For best results use spray mixture the same day it is prepared. Spray uniformly over the area to be treated with a properly calibrated power sprayer, apply as a full coverage spray to run-off when conditions are favorable for disease development. Application should be made to plants when both foliage and flowers are dry, or nearly dry. Spectro can provide both curative and preventive action. Use the highest recommended rate and shortest application interval under conditions of severe disease pressure.

RATES: MAXIMUM INDIVIDUAL APPLICATION RATES

Ornamentals	2.15 lbs/A
Roses	1.53 lbs/A
Pachysandra	
Conifers	
Conifers Seed Beds	

See product label for complete directions for use.

PRECAUTIONS AND RESTRICTIONS

Do not use mistblower or high-pressure spray equipment when making applications in greenhouses.

Do not use this product on plants bearing fruit and other parts or structures that may be eaten.

Do not apply when wind speed favors spray drift beyond the area intended for treatment.

See product label for complete application precautions and restrictions.

