

# QUALI-PRO<sup>®</sup>

# Strobe™ T

## BROAD SPECTRUM FUNGICIDE

FOR CONTROL OF GOLF COURSE TURF AND ORNAMENTAL DISEASES

**Not for use in California**

### ACTIVE INGREDIENT:

Azoxystrobin: methyl (E)-2-[[[6-(2-cyanophenoxy)-4-pyrimidinyl]oxy]-alpha-(methoxymethylene)benzeneacetate..... 11.00%  
Tebuconazole: (+)-alpha-[2-(4-chlorophenyl)ethyl]-alpha-(1,1-dimethylethyl)-1H-1,2,4-triazole-1-ethanol..... 18.35%

**OTHER INGREDIENTS:**..... 70.65%

**TOTAL:** .....100.0%

Suspension concentrate fungicide containing 1.67lb Tebuconazole and 1.00lb Azoxystrobin per gallon.

## CONTENTS: 1 GALLON

GROUP 3 | 11 | FUNGICIDES



**KEEP OUT OF  
REACH OF CHILDREN  
WARNING/AVISO**

See additional precautionary statements  
and directions for use inside booklet.

EPA Reg. No. 53883-358

EPA Est. No. 53883-TX-002

Manufactured by:

**Control  
Solutions Inc.**

5903 Genoa-Red Bluff  
Pasadena, TX 77507



0 72693 00167 1

# QUALI-PRO

## Strobe™ T

### A BROAD SPECTRUM FUNGICIDE

FOR CONTROL OF GOLF COURSE TURF AND ORNAMENTAL DISEASES

Not for use in California



**ACTIVE INGREDIENT:** % BY WT.

Azoxystrobin: methyl E)-2-[[6-(2-cyanophenoxy)-4-pyrimidinyl]oxy]-alpha-(methoxymethylene)benzeneacetate .....	11.00%
Tebuconazole: (+)-alpha-[2-(4-chlorophenyl)ethyl]-alpha-(1,1-dimethylethyl)-1H-1,2,4-triazole-1-ethanol : .....	18.35%
<b>OTHER INGREDIENTS:</b> .....	<b>70.65%</b>
<b>TOTAL:</b> .....	<b>100.00%</b>

Suspension concentrate fungicide containing 1.67lb Tebuconazole and 1.00lb Azoxystrobin per gallon.

## CONTENTS: 1 GALLON

GROUP 3 11 FUNGICIDES

KEEP OUT OF  
REACH OF CHILDREN  
**WARNING/AVISO**

See additional precautionary statements and directions for use inside booklet

EPA Reg. No. 53883-358

EPA Est. No. 53883-TX-002

Manufactured by:



**Control  
Solutions Inc.**

5903 Genoa-Red Bluff  
Pasadena, TX 77507

<b>FIRST AID</b>	
<b>IF SWALLOWED:</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>IF ON SKIN OR CLOTHING:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF IN EYES:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes; then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF INHALED:</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance; then give artificial respiration, preferably mouth-to-mouth if possible.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact SafetyCall® at (866) 897-8050 for emergency medical treatment information.</p>	

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
WARNING**

May be fatal if swallowed. Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing.

**PERSONAL PROTECTIVE EQUIPMENT (PPE)**

**Applicators and other handlers must wear:**

- Coveralls worn over short sleeved shirt and short pants
- Chemical-resistant footwear plus socks
- Chemical-resistant gloves made of any waterproof material (barrier laminate, butyl rubber, neoprene rubber and polyethylene).

**User Safety Requirements:** Follow manufacturer's instructions for cleaning/maintaining personal protective equipment, PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product. Do not reuse them.

### **USER SAFETY RECOMMENDATIONS**

**Users should:**

- Wash hands thoroughly with soap and water before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### **ENVIRONMENTAL HAZARDS**

This pesticide is toxic to mammals, fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

**GROUND WATER ADVISORY:** Azoxystrobin can be persistent for several months or longer. Azoxystrobin has degradation products which have properties similar to chemicals which are known to leach through soil to ground water under certain conditions as a result of agricultural use. Tebuconazole is known to leach through soil into ground under certain conditions as a result of label use. Use of Strobe™ T in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

**SURFACE WATER ADVISORY:** This product may contaminate water through drift or spray in wind. This product has a high potential for runoff into surface waters for several days to weeks after application. Poorly draining soils, wet soils with readily visible slopes toward adjacent surface waters or soil with shallow water tables are more prone to runoff that contains this product. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams and springs will reduce the potential for contamination of water from rainfall runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted within 48 hours. Notify state and/or Federal authorities and Control Solutions, Inc. immediately if you observe any adverse environmental effects due to use of this product.

## **PHYSICAL AND CHEMICAL HAZARDS**

Do not mix or allow coming in contact with oxidizing agent. Hazardous chemical reaction may occur.

## **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read entire label before using this product. This label must be in the possession of the user at the time of pesticide application.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

## **AGRICULTURAL USE REQUIREMENTS**

Use this product in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants, soil, or water is:

- Coveralls worn over short sleeved shirt and short pants
- Chemical-resistant gloves made of any waterproof material (barrier laminate, butyl rubber, neoprene rubber and polyethylene
- Chemical-resistant footwear plus socks

Notify workers of the application by warning them orally and by posting warning signs at entrances to treated areas.

### **NON-AGRICULTURAL USE REQUIREMENTS**

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. The area treated must be vacated by unprotected persons.

**Do not** enter or allow others to enter until sprays have dried.

**Do not** treat areas while unprotected humans or domestic animals are present in the treatment area. Because certain States may require more restrictive reentry intervals, consult your State Department Agriculture for further information

### **OBSERVE THE FOLLOWING RESTRICTIONS WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES, RESERVOIRS, RIVERS, PERMANENT STREAMS, MARSHES OR NATURAL PONDS, AND ESTUARIES.**

- Apply only during alternate years in fields adjacent to aquatic areas listed above.
- Do not apply by ground or air within 100 feet of aquatic areas listed above.
- Do not cultivate within 10 feet of an aquatic area to allow growth of a vegetative filter strip.

### **PRODUCT INFORMATION**

**STROBE™ T** is a suspension concentrate product containing fungicides with multiple modes of action, to provide broad-spectrum preventative and curative control of all major turf diseases that injure cool- and warm-season turf on golf courses. **STROBE™ T** can also be used for disease control in field, nursery and container ornamentals, and commercial and residential landscapes.

### **RESISTANCE MANAGEMENT**

**STROBE™ T** contains both a Group 3 (tebuconazole) and Group 11 (azoxystrobin) fungicide. Fungal diseases may develop resistance to Group 3 and/or Group 11 fungicides when used repeatedly in the same location or in successive years as the primary method of control for targeted diseases. This may result in partial or total loss of control of those species by **STROBE™ T** and/or other Group 3 and/or Group 11 fungicides.

Because the speed and scope of resistant population development cannot be predicted, the use of this product should conform to resistance management strategies established for the crop and use area. Such strategies may include the rotation and/or tank mixing with products utilizing different modes of action or limiting the number of applications per season. Contact your local university or extension specialist and/or manufacturer for fungicide resistance management recommendations.

## SPRAY PREPARATION

Begin application with a clean spray tank (remove scale, pesticide residues, and other foreign matter, and flush system with clean water).

### Mixing Instructions:

1. Fill the tank with 1/2 or 3/4 the desired amount of water.
2. Start mechanical or hydraulic agitation.
3. Slowly add the specified amount of **STROBE™ T**.
4. Add the remaining volume of water.

Maintain continuous agitation to keep the material in suspension. For best results, use spray mixture the same day it is prepared.

## DISEASE CONTROL IN GOLF COURSE TURF

### USE RESTRICTIONS:

- For use on golf course turf only.
- Intended for use by professional applicators.
- Not for residential use.
- Not for use on turf being grown for sale or commercial use as sod.
- Do not use on home lawns and turf sites associated with apartment buildings, daycare centers, playgrounds, playfields etc.
- Do not use clippings for animal feed.
- Do not exceed 7.75 fl. oz. product per 1000 sq. ft. (or 337 fl. oz. product per acre ) (4.4 lb.ai. tebuconazole/A) per year.
- In the state of New York, do not exceed 3.875 fl. oz. of this product per 1,000 sq. ft. (2.2 lb. a.i. tebuconazole/A) per year.
- Do not apply by air or through any type of Ultra Low Volume (ULV) spray system (<5 gallons per acre).
- Do not apply more than 6 applications per year at the lowest rate.

### TURF TYPES:

All cool- and warm-season grasses (such as but not limited to Bentgrasses, Bermudagrasses, Bluegrasses, Fescues, Ryegrasses, St. Augustinegrasses, Zoysiagrasses and Paspalum grasses) or their mixtures.

**CONTROLLING TURF DISEASES:**

**STROBE™ T** can be used for the prevention and control of the diseases mentioned in the table below. Begin applications when conditions favor disease development and repeat applications as long as these conditions persist. Preventative treatments can be applied using 14-21 day intervals as indicated. When treating golf greens, always treat aprons and approaches. Spray uniformly over the area to be treated with properly calibrated equipment. For severe disease pressure use the highest rate and shortest interval listed in the table below.

**TURF APPLICATION INSTRUCTIONS:**

Apply the specified amount of **STROBE™ T** in sufficient water for thorough coverage. Use a volume of 65-130 gallons per acre (1.5 – 3.0 gallons per 1,000 sq. ft.). Apply using properly calibrated low volume, hand held, mechanical or motorized ground broadcast equipment. Application to small areas may be made with low-pressure handwand or backpack equipment. Maintain constant agitation during application. Make all applications after mowing and allow foliage to dry thoroughly before irrigation.

**Note:** Prior to pouring, slowly invert container several times to assure uniform mixture.

**GOLF COURSE TURF APPLICATION INSTRUCTIONS**

<b>DISEASE(S) CONTROLLED</b>	<b>Rate of STROBE™ T fl. oz./1,000 sq. ft.</b>	<b>REMARKS</b>
Anthracnose (Basal** and Foliar) ( <i>Colletotrichum cereale</i> )	0.75 – 1.5	Use preventatively. Begin applications when conditions are favorable for disease infection, prior to disease symptom development.  Repeat application after 14 to 21 days.  For greens a second application may be made after 14 days.  **Use the higher specified rate of 1.5 fl. oz. and shorter intervals of 14 days for Basal Anthracnose.
Bentgrass Dead Spot ( <i>Ophiosphaerella Agrostis</i> )	0.75 – 1.5	Begin applications prior to disease.  Repeat application after 14 days.

(continued)

<b>DISEASE(S) CONTROLLED</b>	<b>Rate of STROBE™ T fl. oz./1,000 sq. ft.</b>	<b>REMARKS</b>
Brown Patch and Zoysia Patch, Large Patch, Leaf and Sheath Spot ( <i>Rhizoctonia solani</i> , <i>R.spp.</i> , <i>R. zeae</i> )	0.75 - 1.5	<p>For Brown Patch. Apply <b>STROBE™ T</b> when conditions are favorable for disease development. Repeat application after 14 to 21 days.</p> <p>For Zoysia Patch, make 1 or 2 applications approximately one month prior to zoysiagrass dormancy. Repeat application after 14 to 21 days.</p> <p>For Large Patch of all warm season turfgrasses, make 1 or 2 applications in the fall prior to infection or when conditions are favorable for infection. Repeat application after 14 to 21 days.</p> <p>For Leaf &amp; Sheath Spot, apply when conditions are favorable for infection such as sequential or periods of temperatures at or above 90 degrees Fahrenheit, Curative control may require several applications. Spray should be targeted at the crown of the turfgrass. Repeat application after 14 to 21 days.</p>
Cool Weather Brown Patch, Yellow Patch ( <i>Rhizoctonia cerealis</i> )	1.5	Make 1 or 2 applications in the fall when conditions are favorable for disease development. Repeat application after 14 to 21 days.
Dollar Spot ( <i>Sclerotinia homoeocarpa</i> )	0.75 – 1.5	Apply preventatively when conditions are favorable for disease development. Repeat application after 14 to 21 days.
Fairy Ring ( <i>Lycoperdon spp.</i> , <i>Agrocybe pediades</i> , and <i>Bovistra plumbea</i> )	0.75 - 1.5	Apply as symptoms develop. Apply in 2-4 gallons of water per 1000 sq. ft. spray volume. Symptoms may take 2-3 weeks to disappear following application. Repeat applications may be required after 28 days in certain cases.

(continued)

<b>DISEASE(S) CONTROLLED</b>	<b>Rate of STROBE™ T fl. oz./1,000 sq. ft.</b>	<b>REMARKS</b>
Gray Leaf Spot ( <i>pyricularia grisea</i> )	0.75 - 1.5	Apply <b>STROBE™ T</b> in a preventative disease control program. Apply before disease is present and alternate with other fungicide chemistries that control gray leaf spot. Repeat application after 14 to 21 days.
Gray Snow Mold	2.4	Apply <b>STROBE™ T</b> once in late fall prior to snow cover.
Pink Snow Mold ( <i>Microdochium nivale</i> ) Fusarium patch (Prior to snow cover) (Intermittent snow cover and rainfall)	2.4 0.75-1.5	Preventative application prior to snow cover. Begin applications when conditions are favorable for disease infection, prior to disease symptom development. Repeat application after 14 to 21 days.
Leaf Rust Stem Rust Stripe Rust ( <i>Puccinia spp.</i> )	0.75 – 1.5	Begin applications when conditions are favorable for disease infection, prior to disease symptom development Repeat application after 14 to 21 days.
Leaf Spot ( <i>Bipolaris sorokiniana</i> )	0.75 – 1.5	Apply when conditions are favorable for disease development. Repeat application after 14 to 21 days.
Melting Out ( <i>Dechslera poae</i> )	0.75 – 1.5	Apply when conditions are favorable for disease development. Repeat application after 14 to 21 days.
Necrotic Ring Spot ( <i>Ophiosphaerella korrae</i> )	1.5	Apply when conditions are favorable for disease development. Repeat application after 14 to 21 days.

(continued)

<b>DISEASE(S) CONTROLLED</b>	<b>Rate of STROBE™ T fl. oz./1,000 sq. ft.</b>	<b>REMARKS</b>
Pink Patch ( <i>Limonomyses roseipellis</i> )	0.75 – 1.5	Apply when conditions are favorable for disease development. Repeat application after 14 to 21 days.
Powdery Mildew ( <i>Erysiphe graminis</i> )	0.75 – 1.5	Begin applications when conditions are favorable for disease infection, prior to disease symptom development. Repeat application after 14 to 21 days.
Pythium Blight Pythium Root Rot ( <i>Pythium aphanidermatum</i> , <i>Pythium spp.</i> )	0.75 – 1.5	Use preventatively. Begin applications when conditions are favorable for disease infection, prior to disease symptom development. During periods of prolonged favorable conditions, treat on the 10 day application interval. For use on newly seeded as well as established turf. Repeat application after 14 to 21 days.
Red Thread ( <i>Laetisaria fuciformis</i> )	0.75 – 1.5	Apply when conditions are favorable for disease development. Repeat application after 14 to 21 days.
Southern Blight ( <i>Sclerotium rolfsii</i> )	0.75 – 1.5	Apply when conditions are favorable for disease development. Repeat application after 14 to 21 days.
Spring Dead Spot ( <i>Ophiosphaerella korrae</i> ) or ( <i>Ophiosphaerella narmari</i> ) or ( <i>Ophiosphaerella herpotricha</i> )	1.5	Apply 1 or 2 applications approximately one month prior to bermudagrass dormancy. Irrigate ¼" to ½." directly after application. Repeat application after 14 to 21 days.

## DISEASE CONTROL IN FIELD, NURSERY AND CONTAINER ORNAMENTALS AND COMMERCIAL and RESIDENTIAL LANDSCAPE ORNAMENTALS

### USE RESTRICTIONS

- Intended for use by professional applicators
- Do not apply to bearing fruit trees or vegetables
- Do not use clippings for animal feed
- Do not exceed 1.84 fl oz per 1000 sq ft per year (80 fl oz per acre per year).
- Do not exceed the maximum single application rate of 20 fl. oz. of **STROBE™ T** per acre or the maximum application rate of 80 fl. oz. per acre per year.
- Do not exceed 4 applications per year at the highest specified rate.

ORNAMENTAL PLANT	
Herbaceous Bedding	Ageratum, Begonia, Canna, Coleus, Dahlia, Dusty Miller, Foxglove, Fuchsia, Geranium, Impatiens, Lavender, Marigold, Pansy, Petunia, Pinks, Primrose, Salvia, Statice, Strawflower, Tickseed, Verbena
Flowering	Chrysanthemum, Hydrangea, Hollyhock, Iris, Lily, Poinsettia
Tropical Foliage	Dieffenbachia, Dracaena, English Ivy, Philodendron, Pothos
Woody Ornamentals	Azalea, Hibiscus, Holly, Ligustrum, Rhodendrum, Rose, Pyracantha
Evergreen Trees	Douglas Fir, Fir, Larch, Pine, Spruce
Deciduous Trees	Ash, London Plane, Maple, Oak, Sycamore, Walnut
Flowering Trees	Cherry, Crabapple, Hawthorn, Mountain Ash, Pear

## CONTROLLING ORNAMENTAL DISEASES

**STROBE™ T** provides broad-spectrum control of foliar, stem, and below-ground diseases on ornamental plants. Apply **STROBE™ T** prior to disease development and at the very latest, upon first sign of disease.

### ORNAMENTALS AND DISEASES CONTROLLED

ORNAMENTAL TYPES	DISEASE(S)
Woody ornamentals, shade trees	Anthracoze
Herbaceous ornamentals	Ascochyta Blight
Roses	Black spot
Woody and herbaceous ornamentals	Botrytris (Gray Mold)
Woody and herbaceous ornamentals	Brown Rot
Woody and herbaceous ornamentals	Colletotrichum
Woody and herbaceous ornamentals	Cercospora Leaf Spot
Ligustrum	Corynespora Leaf Spot
Iris	Didymellina Leaf Spot
Shade and ornamental trees	Diplodia Tip Blight (Diplodia pinea)
Azalea, rhododendron <sup>2</sup>	Ovulinia
Woody and herbaceous ornamentals	Entomosporium Leaf Spot
Woody and herbaceous ornamentals	Fusicaladium Leaf Scab
Woody and herbaceous ornamentals	Phomopsis Blight
Woody ornamentals	Pine Tip Blights
Woody and herbaceous ornamentals, ornamental nut and fruit trees	Powdery Mildew
Ornamental nut and fruit trees	Rust Diseases
Herbaceous ornamentals	Ramularia Leaf Spot
Pyracantha, flowering crab, ornamental fruit and nut trees	Scab
Woody and herbaceous ornamentals	Septoria Leaf Spot
Woody and herbaceous ornamentals	Venturia Leaf Scab

**Note: STROBE™ T** has been determined to be safe for use on the plant types listed in these directions for use based on cumulative data derived from research trials and historical field use. As all species and cultivars have not been tested, perform trial applications if a user wishes to make an application to a plant type not listed on the label but found on a similar use site and for disease that is listed on the label. To conduct a trial application, apply at least two applications to at least 25 trial plants at the highest concentration, 7 days apart. Evaluate 7 days after the last application before initiating full-scale application. Do not use this product on the following plants: Swedish Ivy (*Nephrrolepis exaltata*), Boston fern (*Plectranthus australis*), and Easter cactus (*Hatiora gaertneri*).

#### ORNAMENTAL APPLICATION INSTRUCTIONS

Apply **STROBE™ T** with properly calibrated hand held, mechanical or motorized spray equipment. Use the specified amount of **STROBE™ T** Fungicide per 100 gallons of water or adequate spray volume to obtain uniform coverage. Begin applications when disease first appears and repeat at 14-21 day intervals during the growing season. Use the shortest interval when conditions are usually favorable for the development of coverage spray to drip volume appropriate for the size of the plants and amount of foliage, which will provide thorough coverage throughout the canopy. Allow sprays to dry before overhead irrigation is applied.

DISEASE(S) CONTROLLED	RATE OF STROBE™ T fl. oz./100 Gallons/A or Pints/A	REMARKS
Anthraxnose ( <i>Colletotrichum</i> )	5- 10 fl. oz. per 100 gallons of water or 0.625-1.25 pts/A	Apply as buds break or at first sign of disease. Repeat application at 14-21 day intervals as needed during disease period.
Black Spot of Rose ( <i>Diplocarpon rosae</i> )	5- 10 fl. oz. per 100 gallons of water or 0.625-1.25 pts/A	Apply early summer or at first sign of disease. Repeat application every 14-21 days as needed during disease period.
Brown Rot and Blight ( <i>Monilinia</i> , <i>Sclerotinia</i> , <i>Whetzelinia</i> )	5- 10 fl. oz. per 100 gallons of water or 0.625-1.25 pts/A	Apply late spring or at first sign of disease. Repeat application every 14-21 days as needed during the disease period.
Fusicladium and Venturia Leaf Scabs on: Crabapple, Hawthorn, Pear, Mountain Ash, Pyracantha, etc.	5- 10 fl. oz. per 100 gallons of water or 0.625-1.25 pts/A	Apply as buds break. Repeat application every 14-21 days during disease period. Effective control requires coverage during expansion. Rotations with strobilurin (QoI) fungicides can be utilized.

(continued)

DISEASE(S) CONTROLLED	RATE OF STROBE™ T fl. oz./100 Gallons/A or Pints/A	REMARKS
Leaf Spots and Blights caused by: <i>Ascochyta</i> , <i>Blumeriella</i> , <i>Botrytis</i> , <i>Cercospora</i> , <i>Coccomyces</i> , <i>Corynespora</i> , <i>Curvularia</i> , <i>Didymellina</i> , <i>Entomosporium</i> , <i>Fabraea</i> , <i>Fusarium</i> , <i>Ramularia</i> , <i>Rhizoctonia</i> , <i>Marssoninia</i> , <i>Mycosphaerella</i> , <i>Myrothecium</i> , <i>Phoma</i> , <i>Physalasporea</i> , <i>Schizothyrium</i> , <i>Septoria</i> , <i>Sphaceloma</i>	5- 10 fl. oz. per 100 gallons of water or 0.625-1.25 pts/A	Make applications when disease symptoms first appear. Repeat every 14-21 days during disease period. Rotations with strobilurin (QoI) can be used.
Ovulinia Blight	5- 10 fl. oz. per 100 gallons of water or 0.625-1.25 pts/A	Apply as flowers open. Repeat every 14 to 21 days during disease period.
Powdery Mildews <i>Erysiphe</i> , <i>Microsphaera</i> , <i>Phyllactinia</i> , <i>Podosphaera</i> , <i>Oidium</i> , <i>Sphaerotheca</i>	5- 10 fl. oz per 100 gallons of water or 0.625-1.25 pts/A	Apply when disease first appears and repeat application every 14 to 21 days. Rotations with other effective products can be used.
Rust Diseases caused by: <i>Puccinia</i> , <i>Gymnosporangium</i> , <i>Uromyces</i>	5- 10 fl. oz per 100 gallons of water or 0.625-1.25 pts/A	Apply late spring or when symptoms first appear. Repeat applications every 14 to 21 days during disease period. Rotations with other effective products can be used.
Tip Blight of Pine <i>Sphaeropsis sapinea</i> , <i>Diplodia pineae</i>	5- 10 fl. oz per 100 gallons of water or 0.625-1.25 pts/A	Begin applications in the spring when new growth starts. Make a second application just prior to needle emergence from the sheath and a third application 14-21 days later. Ensure thorough coverage.

(continued)

DISEASE(S) CONTROLLED	RATE OF STROBE™ T fl. oz./100 Gallons/A or Pints/A	REMARKS
Twig Blights, Cankers, and Diebacks <i>Diaporthe, Kabatina, Phoma, Phomopsis</i>	5- 10 fl. oz per 100 gallons of water or 0.625-1.25 pts/A	Apply when symptoms first appear. Repeat applications every 14 to 21 days during disease period.

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal or cleaning of equipment.

**PESTICIDE STORAGE:** Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food and feed. Store in original container and out of reach of children, preferably in a locked storage area. Do not store above 100 degrees F for extended period of time. Storage below 20 degrees F can result in formation of crystals. If product crystallizes, store at 50 degrees F to 70 degrees F and agitate to redissolve crystals. If container is damaged or spill occurs, use product immediately or dispose of product and damaged container as indicated below.

**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

### CONTAINER HANDLING:

**Nonrefillable Container (five gallons or less):** Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling if available or, puncture and dispose of in a sanitary landfill or by incineration or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

## LIMITATION OF WARRANTY AND LIABILITY

Read the entire directions for use, conditions of warranties and limitations of liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following **CONDITIONS, DISCLAIMER OF WARRANTIES and LIMITATIONS OF LIABILITY.**

**CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Control Solutions, Inc. All such risks shall be assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** To the extent consistent with applicable law, Control Solutions, Inc. makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond the statements made on this label. No agent of Control Solutions, Inc. is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, Control Solutions, Inc. disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product.

**LIMITATIONS OF LIABILITY:** To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid or at Control Solutions, Inc.'s election, the replacement of product.

Quali-Pro® is a registered trademark of The ADAMA Group

Manufactured by:  
**Control Solutions Inc.**  
Pasadena, Texas 77507

**QUALI-PRO**

**Strobe™ T**

**A BROAD SPECTRUM FUNGICIDE**

**FOR CONTROL OF GOLF COURSE TURF AND ORNAMENTAL DISEASES**

Not for use in California

**ACTIVE INGREDIENT:**

**% BY WT.**

Azoxystrobin: methyl E)-2-[[6-(2-cyanophenoxy)-4-pyrimidinyl]oxy]-alpha-(methoxymethylene) benzeneacetate .....	11.00%
Tebuconazole: (+)-alpha-[2-(4-chlorophenyl)ethyl]-alpha-(1,1-dimethylethyl)-1H-1,2,4-triazole-1-ethanol : .....	18.35%
<b>OTHER INGREDIENTS:</b> .....	<b>70.65%</b>
<b>TOTAL:</b> .....	<b>100.00%</b>

Suspension concentrate fungicide containing 1.67lb Tebuconazole and 1.00lb Azoxystrobin per gallon.

**CONTENTS: 1 GALLON**

GROUP 3 11 FUNGICIDES



**KEEP OUT OF  
REACH OF CHILDREN  
WARNING/AVISO**

See additional precautionary statements and directions for use inside booklet.

EPA Reg. No. 53883-358

EPA Est. No. 53883-TX-002



Manufactured by:

**Control  
Solutions Inc.**

5903 Genoa-Red Bluff  
Pasadena, TX 77507