Supplemental Label





SUREGUARD® HERBICIDE FOR USE IN DORMANT WARM-SEASON TURFGRASS GROWN ON RESIDENTIAL SITES, GOLF COURSES, SOD PRODUCTION AND SIMILAR AREAS

(Only for use in the following states: Alabama, Arizona, Arkansas, Colorado, Delaware, Florida, Georgia, Iowa, Indiana, Illinois, Kansas, Kentucky, Louisiana, Maryland, Mississippi, Missouri, Nebraska, Nevada, New Mexico, New Jersey, North Carolina, Oklahoma, Ohio, Pennsylvania, South Carolina, Tennessee, Texas, Virginia and West Virginia)

This supplemental label expires July 30, 2014 and must not be used or distributed after this date.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF APPLICATION. READ THE LABEL AFFIXED TO THE CONTAINER FOR SUREGAURD HERBICIDE BEFORE APPLYING. USE OF SUREGUARD HERBICIDE ACCORDING TO THIS LABELING IS SUBJECT TO THE USE PRECAUTIONS AND LIMITATIONS IMPOSED BY THE LABEL AFFIXED TO THE CONTAINER FOR SUREGUARD HERBICIDE

DIRECTIONS FOR USE ON DORMANTWARM-SEASON TURFGRASS GROWN ON RESIDENTIAL SITES, GOLF COURSES, SOD PRODUCTION AND SIMILAR AREAS

SureGuard Herbicide may be applied as a single or split application to well established dormant turfgrass listed in Table A, and will control winter annual weeds found in Table 1 (on the container label). Apply SureGuard Herbicide to dormant turfgrass in such areas as apartment complexes, golf courses, sod farms, roadsides, sports fields, campgrounds, office complexes, parks, parking areas, recreational sites, schools and other similar sites. Dormant bermudagrass has exhibited tolerance to SureGuard Herbicide only when applied after turf has become dormant in the late fall and before turf breaks dormancy in the late winter/early spring. Application of SureGuard Herbicide to actively growing turfgrass (warm season and cool season) or during green-up will cause unacceptable injury. SureGuard Herbicide will injure warm season turf grown in southern areas where grass does not become completely dormant.

BROADCAST APPLICATIONS

Apply 8 to 12 oz of *SureGuard* Herbicide per broadcast acre as a preemergence (to weed emergence) application. If weeds are present at the time of application apply *SureGuard* Herbicide plus an adjuvant (0.25% v/v non-ionic surfactant). Make postemergence (to weed emergence) applications of *SureGuard* Herbicide when weeds are actively growing and no larger than 2 inches in height. Thorough spray coverage is necessary to maximize the postemergence activity of *SureGuard* Herbicide. When applied after weed germination, *SureGuard* Herbicide will provide preemergence and postemergence control of broadleaf weeds and grasses listed in Table 1 (on the container label). Postemergence control of *SureGuard* Herbicide may be more effective on certain weed species, and may not control mature, stressed or hardened off weeds that are not actively growing at the time of application.

A second application of *SureGuard* Herbicide may be required to provide adequate season-long weed control. Apply the second application using the above mentioned rate guidelines prior to the turfgrass breaking spring dormancy.

SPOT TREATMENTS

Mix 2 1/2 tsp per gal of *SureGuard* Herbicide and 2 tsp (1/3 fl oz) of non-ionic surfactant in one gal of water and apply one gal of spray solution per 1,000 sq ft. Occasionally shake the spray solution while spraying to ensure the spray solution remains well mixed. Spray the target weeds until the leaves are wet.

TANK MIXING WITH OTHER TURFGRASS HERBICIDES

Tank mixing *SureGuard* Herbicide with other preemergence and postemergence herbicides registered for use in dormant turfgrass may provide a broader spectrum of weed control than *SureGuard* Herbicide alone.

IMPORTANT: Turfgrass must be completely dormant at application. Any turfgrass that is not dormant will be injured by applications of *SureGuard* Herbicide. Scout area to be sprayed for any turf that is green in color and if encountered, delay application until turfgrass is completely dormant. Read and follow the label of any herbicides mixed with *SureGuard* Herbicide. When tank mixing *SureGuard* Herbicide with other herbicides, always follow the most restrictive limitations and precautions on the label of any tank mix partner.

USE PRECAUTIONS

Exercise good judgment and caution when applying to dormant turfgrass until familiarity is gained with *SureGuard* Herbicide.

RESTRICTIONS AND LIMITATIONS

- Do not apply to golf course putting greens.
- Do not apply to warm season turfgrass that has been overseeded with cool season turfgrass (ex. perennial rye).
- Do not irrigate within 1 hour before or after application.
- Do not apply if rain is expected within 1 hour after application.
- Do not mow turfgrass within 12 hours after application.
- Do not apply within 30 days prior to cutting or lifting sod.
- Do not apply more than 2 applications at 12 oz/A or 3 applications at 8 oz/A per year.
- Do not re-apply SureGuard Herbicide within 30 days.
- Do not apply in fall before turfgrass has ceased active growth or in late winter/early spring after turfgrass has resumed active growth.
- Allow 8 weeks between application and seeding or sodding of turfgrass.

	TABLE A:	TOLERANT	TURFGRASS SPECIES
--	----------	----------	-------------------

COMMON NAME	SCIENTIFIC NAME
Bermudagrass	Cynodon spp.

PLEASE CONTACT VALENT U.S.A. CORPORATION AT 800-6-VALENT (682-5368) TO DETERMINE IF THIS USE IS REGISTERED IN YOUR STATE.

Copyright © 2011 by Valent U.S.A. Corporation

SureGuard and Products That Work, From People Who Care are registered trademarks of Valent U.S.A. Corporation

Manufactured for: **Valent U.S.A. Corporation** P. O. Box 8025 Walnut Creek, CA 94596-8025

Made in U.S.A.

2011-SRG-0010 20110401

Products That Work, From People Who Care® | www.valentpro.com | 800-89-VALENT (898-2536)