



Surflan[®] XL 2G
HERBICIDE

THE CHOICE FOR LANDSCAPE PROFESSIONALS SEEKING EFFECTIVE, ECONOMICAL PRE-EMERGENT WEED CONTROL

HRAC Class: Group 3 - herbicide

Active ingredient: *Benefin + oryzalin*

Formulation: granular

Package size: 50 lb bag

APPLICATION USAGE

Established tall fescue and warm season turf (bahagrass, Bermudagrass, buffalograss, centipedegrass, St. Augustinegrass, and zoysiagrass)	Established flowers Non-bearing fruit and nut trees, non-bearing vineyards and non-bearing berries Christmas tree plantations Landscape ornamentals Ground covers
Container grown ornamentals	Ornamental bulbs Non-cropland and
Nursery stock	bareground

APPLICATION RATES*

Ornamentals	2.3 - 3.4 lbs/1,000 sq ft
Ornamental bulbs	4.6 - 6.9 lbs/1,000 sq ft
Warm season turf	1.7 - 3.4 lbs/1,000 sq ft

*Refer to product label for specific use rates.

Surflan[®] XL 2G herbicide stops unsightly weeds before they get started. Surflan XL 2G is an easy-to-use selective pre-emergence herbicide for control of certain annual grasses and broadleaf weeds. Used according to label directions, Surflan XL 2G can provide effective weed control all season long.

Apply Surflan XL 2G prior to germination of target weeds, or immediately after cultivation. Length of weed control will vary depending upon the rate of Surflan XL 2G applied, weed population, soil conditions, temperature, watering regime and other factors. Following application, user should monitor and observe level of weed control over time to determine if additional applications may be needed.

- **Weeds controlled**

Labeled to control or suppress 21 annual grasses and 35 broadleaf weeds. See label for details.

- **Plant species labeled for application**

Surflan XL 2G is labeled for use on over 250 species of trees, shrubs, perennials, annuals. See label for details.

- **Safe for ornamental bulbs**

See label for additional information.

- **Label flexibility**

Users who wish to use Surflan XL 2G on plant species not recommended on this label may determine the suitability for such uses by treating a small number of such plants at a recommended rate. Prior to treatment of larger areas, the treated plants should be observed for any sign of herbicidal injury during 30 to 60 days of normal growing conditions to determine if the treatment is non-injurious to the target plant species. The user assumes responsibility for any plant damage or other liability resulting from the use of Surflan XL 2G on plant species not recommended on the label.



Contact your local UPI sales representative or distributor, call 1-800-438-6071 or visit www.upi-usa.com to learn more.

Always read and follow label directions and precautions. Surflan[®] and UPI logo are registered trademarks of United Phosphorus, Inc. ©July, 2017. UPI, 630 Freedom Business Center, King of Prussia, PA 19406.

SURFLAN XL 2G - SPD - 17 - 01