



Trigo™

Product information sheet

// Active Ingredients: Trifloxystrobin 8.33%
Triadimefon 41.67%

// IRAC Group: 11, 3


// EPA Registration Number: 432-1513

// Signal Word: Caution

// Formulation: Wettable Dispersible Granule

// REI: 12 hours

Trigo™ fungicide combines the strength of two active ingredients to maximize disease control with long-lasting residual activity.



Best Uses

// Control of 24 pathogens including key foliar diseases: Anthracnose, Black Spot, Powdery Mildew, Rust, Scab and leaf spot diseases caused by *Cercospora*, *Entomosporium* and *Septoria*

// In rotation, as this unique combination of active ingredients is ideal for resistance management

Attributes

// Up to 30 days of disease control including new growth

// Mesostemic (vapor barrier) and translaminar movement provides disease control in and around the plant tissue

// Use on nursery and greenhouse ornamentals


Application Rates

// For greenhouse applications, apply 1.2-2.4 oz./100 gal. of water


// For nursery applications, apply 3-9 oz./100 gal. of water

// Trigo is available in a 1-lb. package

Rust Control on Plumeria



Plumeria Rust (*Coleosporium plumeriae*)



(Aaron Palmateer, Bayer)

(Aaron Palmateer, Bayer)

backedbybayer.com | backedbybayer.com/production-ornamentals



ALWAYS READ AND FOLLOW LABEL INSTRUCTIONS
 Bayer CropScience LP, Environmental Science Division, 2 T.W. Alexander Drive, Research Triangle Park, NC 27709. For additional product information, call toll-free 1-800-331-2867. Not all products are registered in all states. Bayer (reg'd), the Bayer Cross (reg'd), and Trigo™ are trademarks of Bayer. Pageant is a registered trademark of BASF. © 2018 Bayer CropScience LP. ES-418-TRG-144-A-R1