

[CLICK HERE TO JUMP TO USE DIRECTIONS](#)

GORDON'S® TRIMEC® CRABGRASS PLUS LAWN WEED KILLER READY-TO-USE

*Also Kills Dandelion & Clover
Plus 200 Other Listed Weeds*

ACTIVE INGREDIENTS:

2,4-D, dimethylamine salt	0.313%
Quinclorac	0.104%
Dicamba, dimethylamine salt	0.029%
OTHER INGREDIENTS:	99.554%

TOTAL 100.000%

THIS PRODUCT CONTAINS:

2,4-dichlorophenoxyacetic acid equivalent, 0.260%
3,7-dichloro-8-quinolinecarboxylic acid, 0.104%
3,6-dichloro-o-anisic acid equivalent, 0.024%
Isomer Specific by AOAC Method.

KEEP OUT OF REACH OF CHILDREN CAUTION



READ THE ENTIRE LABEL FIRST.
OBSERVE ALL PRECAUTIONS AND
FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Causes moderate eye irritation. Harmful if swallowed. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco. Remove and wash clothing before reuse.

Applicators and other handlers of this product must wear:

- Long sleeved shirt and long pants
- Shoes
- Socks

First Aid	
If in eyes:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to by a poison control center or doctor.• Do not give anything to an unconscious person.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment advice.	

Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

GENERAL PRECAUTIONS AND RESTRICTIONS:

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until sprays have dried.
- Apply when daytime temperatures are between 50° and 90°F (do not apply above 85°F for bermudagrass). Do not treat when air temperatures exceed 90°F as damage to turfgrass may occur.
- Do not apply this product to food crops (gardens, fruits and vegetables), forage crops, or ornamental plants including flowers, trees, shrubs, hedges, woody ornamentals, groundcovers established in landscape plantings and other nontarget plants.
- Do not spray exposed roots of ornamentals and trees.

SHAKE WELL BEFORE USING

1. WHAT IT DOES: Kills major broadleaf and troublesome grass weeds including dandelion, chickweed, clover as well as crabgrass and foxtail.

2. HOW IT WORKS: Trimec® Crabgrass Plus Lawn Weed Killer Ready-To-Use contains three proven weed killers that target lawn weeds and crabgrass. The herbicides enter the lawn weeds through their leaves and move inside the weeds, even down into the roots. And lawns listed on this label will not be harmed when used as directed.

3. WHERE TO USE: Trimec Crabgrass Plus Lawn Weed Killer Ready-To-Use is a ready-to-use (RTU) product intended for the treatment of individual weeds, patches of weeds or sparse weeds scattered throughout home lawns. Trimec Crabgrass Plus Lawn Weed Killer Ready-To-Use can be applied to the following established turfgrass species:

Cool-season turfgrass

Kentucky bluegrass, Ryegrass (perennial and annual), Tall fescue, Mixtures of cool season grasses containing fine fescues (red, hard and chewings) and bentgrass.

Warm-season turfgrass

Bermudagrass (common and hybrid), Zoysiagrass, Buffalograss

DO NOT apply Trimec Crabgrass Plus Lawn Weed Killer Ready-To-Use to the following:

Warm-season turfgrass

St. Augustinegrass, Bahiagrass, Centipedegrass, Carpetgrass, Seashore paspalum

Other areas

Lawns with desirable clovers or legumes, Dichondra, Ornamental plants (flowers, trees, groundcovers, landscape beds and shrubs), Turfgrass species that are not listed on this label. Gardens including vegetables, fruit trees, vines and berries.

4. WEEDS CONTROLLED

GRASS WEEDS:

crabgrass (large and smooth)	foxtail (green, yellow and giant)	kikuyugrass broadleaf signalgrass
------------------------------	-----------------------------------	--------------------------------------

BROADLEAF WEEDS: A partial list is presented below:

Annual yellow	Bull thistle	Cudweed
sweetclover	Bullnettle	Curly dock
Aster	Burclover	Daisy, English
Austrian fieldcress	Burdock	Daisy fleabane
Bedstraw	Burning nettle	Daisy, oxeye
Beggarticks	Burweed	Dandelion
Betony, Florida	Buttercup	Dichondra
Bindweed, field	Buttonweed	Dogbane
Bird vetch	Canada thistle	Dogfennel
Bitter wintercress	Carolina geranium	Dollarweed
Bittercress, hairy	Carpetweed	Elderberry
Bitterweed	Catchweed bedstraw	False dandelion
Black-eyed Susan	Catnip	Falseflax
Black medic	Catsear	False sunflower
Black mustard	Chickweed, common	Fiddleneck
Blackseed plantain	Chickweed, mouseear	Florida pusley
Blessed thistle	Chicory	Frenchweed
Bloodflower milkweed	Cinquefoil	Galinsoga
Blue lettuce	Clover, crimson	Goathead
Blue vervain	Clover, hop	Goldenrod
Bracted plantain	Clover, red	Ground ivy
Brassbuttons	Clover, strawberry	Gumweed
Bristly oxtongue	Clover, sweet	Hairy fleabane
Broadleaf dock	Clover, white	Hawkweed
Broadleaf plantain	Cockle	Healall
Broomweed	Cocklebur	Heartleaf drymary
Buckhorn	Common mullein	Heathaster
Buckhorn plantain	Creeping Jenny	Hedge bindweed
Bulbous buttercup	(Creeping Charlie)	(continued on next page)

BROADLEAF WEEDS (cont.):

Hedge mustard	Poorjoe	Thistle
Hemp	Prairie sunflower	Trailing crownvetch
Henbit	Prickly lettuce	Tumble mustard
Hoary cress	Prickly sida	Tumble pigweed
Hoary plantain	Prostrate knotweed	Velvetleaf
Hoary vervain	Prostrate pigweed	Venice mallow
Horsenettle	Prostrate spurge	Virginia buttonweed
Jimsonweed	Prostrate vervain	Virginia creeper
Knawel	Puncturevine	Virginia pepperweed
Knotweed	Purslane, common	Water pennywort
Kochia	Ragweed	Wavyleaf bullthistle
Lambsquarters	Red sorrel	Western clematis
Lespedeza	Redroot pigweed	Western salsify
Mallow	Redstem filaree	White mustard
Matchweed	Rough cinquefoil	Wild aster
Mexicanweed	Rough fleabane	Wild buckwheat
Milk vetch	Russian pigweed	Wild carrot
Morningglory	Russian thistle	Wild four-o'clock
Mouseear hawkweed	Scarlet pimpernel	Wild garlic
Mugwort	Scotch thistle	Wild geranium
Musk thistle	Sheep sorrel	Wild lettuce
Mustard	Shepherdspurse	Wild marigold
Narrowleaf plantain	Slender plantain	Wild mustard
Narrowleaf vetch	Smallflower galinsoga	Wild onion
Nettle	Smooth dock	Wild parsnip
Orange hawkweed	Smooth pigweed	Wild radish
Oriental cocklebur	Sorrel	Wild rape
Oxalis	Sowthistle	Wild strawberry
Parsley-piert	Spanishneedles	Wild sweet potato
Parsnip	Speedwell	Wild vetch
Pearlwort	Spiny amaranth	Wild violet
Pennycress	Spiny cocklebur	Woodsorrel
Pennywort	Spiny sowthistle	Woolly croton
Peppergrass	Spotted catsear	Woolly morningglory
Pepperweed	Spotted spurge	Woolly plantain
Pigweed	Spurweed	Wormseed
Pineywoods bedstraw	Stinging nettle	Yarrow
Plains coreopsis	Strawberry, India mock	Yellow rocket
(tickseed)	Tall nettle	Yellowflower
Plantain	Tall vervain	pepperweed
Poison ivy	Tansy ragwort	and other
Poison oak	Tansy mustard	broadleaf weeds
Pokeweed	Tanweed	

- Avoid applications of this product when turfgrasses are under stress since injury may result.

Mowing:

- Delay mowing 2 days before and until 2 days after the application of this product.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Keep from freezing. Store in original container in a locked storage area inaccessible to children and pets.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED. THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

TRIMEC® is a registered trademark of PBI/Gordon Corporation.

755/8-2009 AP081009
EPA REG. NO. 2217-906



pbi / Gordon
corporation

An Employee-Owned Company

MANUFACTURED BY
PBI/GORDON CORPORATION
1217 WEST 12TH STREET
KANSAS CITY, MISSOURI 64101
www.pbigordon.com

5. WHEN TO USE:

Apply this product when weeds are small and actively growing.

CRABGRASS (large and smooth):

- **Spring:** Apply this product from crabgrass emergence until the crabgrass is 3 to 4 inches in height. This period begins the first time that you can see crabgrass in your lawn. Re-treat at 14 days later for dense populations.

FOXTAIL (GREEN, GIANT AND YELLOW):

- **Spring and early summer:** Apply this product to foxtail during the early growth stages or when foxtail is 1 to 4 inches in height. Re-treat at 14 days later for dense populations.

BROADLEAF WEEDS:

- **Spring:** Apply when dandelion, plantain, buttercup, speedwell, knotweed and other broadleaf weeds are young and actively growing with adequate soil moisture for best results.
- **Fall:** Apply when henbit, chickweed, white clover, ground ivy, lawn burweed and other broadleaf weeds are young and actively growing with adequate soil moisture for best results.

6. HOW TO USE

1. Shake well.
2. Adjust the nozzle to coarse spray pattern to minimize wind drift. Spray the weeds only. Spray to lightly cover broadleaf and grass weeds. Excessive spraying can injure some lawns. Do not apply when daytime temperatures are above 90°F as lawn injury can occur (do not apply above 85°F for bermudagrass).

7. CULTURAL TIPS:

For newly seeded areas:

- Do not apply within 4 weeks after seedling emergence of Kentucky bluegrass, fine fescue mixtures and perennial ryegrass. Apply this product to newly seeded grasses when well-established (approximately 6 weeks after seedling emergence) or after the third mowing.

For newly sodded, sprigged, or plugged areas:

- The application of this product to newly sodded, sprigged, or plugged grasses should be delayed until 3 to 4 weeks after the sodding, sprigging, or plugging operations.

Seeding:

- Turf species listed on this label can be seeded into the treated areas at four (4) weeks after the application of this product.

Irrigation:

- Do not apply this product immediately before rainfall or irrigation. For best results, do not irrigate or water the turfgrass within 24 hours after application.