



**TEBUCONAZOLE 3.6 SELECT CONTAINS TEBUCONAZOLE, THE ACTIVE INGREDIENT  
USED IN TORQUE™**

# **TEBUCONAZOLE 3.6 SELECT™**

FUNGICIDE

## ***Disease Control for Golf Turf and Ornamentals***



### **ACTIVE INGREDIENT**

Tebuconazole

### **% ACTIVE INGREDIENT**

38.7%

### **FORMULATION**

Suspension Concentrate

### **CHEMICAL FAMILY**

Triazole

### **BRAND ALTERNATIVE**

Torque™

### **MODE OF ACTION**

Demethylation Inhibitor

### **RESTRICTED USE**

None

### **SIGNAL WORD**

Caution

### **PACKAGING INFORMATION**

4 x 1 gal.

### **LABELED USES**

Commercial Landscape  
Ornamentals  
Golf Course Turf  
Nursery Ornamentals  
Residential Landscape  
Ornamentals

**Tebuconazole 3.6 Select™** provides preventive and curative control against some of the most difficult diseases. Tebuconazole 3.6 Select can protect your ornamentals for up to 3 weeks against diseases such as mildews, rusts and blights. Tebuconazole 3.6 Select can be used to combat numerous diseases on all golf turf applications of cool and warm season grasses or their mixtures.

- Preventive and curative properties
- Provides control for up to 3 weeks
- Acropetal (upward) movement
- Low use rates
- Rapid uptake (4 hours)
- Highly compatible tank mix partner

**IMPORTANT:  
ALWAYS READ AND FOLLOW  
LABEL INSTRUCTIONS.**

  
**PRIME SOURCE**

# TEBUCONAZOLE 3.6 SELECT™

FUNGICIDE

## **Turf Applications:**

For use on all golf turf applications of cool season and warm season grasses (such as Bentgrasses, Bluegrasses, Fescues, Ryegrasses, St. Augustine grasses, and Zoysia) or their mixtures. Tebuconazole 3.6 Select™ is not phytotoxic to any of the above mentioned grasses when used in accordance with the label.

Note: Bermudagrass can be sensitive to Tebuconazole 3.6 Select under certain conditions. Do not apply consecutive applications during or just after dormancy break. Avoid applications when temperatures are expected to exceed 85°F.

## **Restrictions:**

- For use on golf course turf only.
- Do not use on home lawns and turf sites associated with apartment buildings, daycare centers, playgrounds, playfields, recreational park athletic fields, athletic fields located on or next to schools (i.e., elementary, middle and high school), campgrounds, churches, and theme parks.
- Not for homeowner use.
- Not for use on turf being grown for sale or commercial use as sod.
- Do not use clippings for animal feed.
- Do not exceed 3.6 fl. oz. of Tebuconazole 3.6 Select per 1,000 sq. ft. per year.
- Do not apply more than 6 applications per year except in New York. In New York State, do not apply more than 3 applications per year.

## **Field, Nursery and Container Ornamentals and Commercial and Residential Landscape Applications:**

Tebuconazole 3.6 Select can be used in a preventative and curative disease control program for the listed plant types and disease in the table below. Optimum disease management is obtained when Tebuconazole 3.6 Select is used in conjunction with sound disease management practices. Apply material with properly calibrated hand held, mechanical or motorized spray equipment. Begin applications when disease first appears and repeat at 14-21 day intervals during the growing season. Use the shortest interval when conditions are unusually favorable for the development of disease.

## **Restrictions:**

- For use on ornamental plants only; not for use on woodlands or forest management.
- Not for homeowner use.
- Do not apply more than 10 fl. oz. per acre in a single application.
- Do not apply more than 0.31 gallons (40 fl. oz.) of Tebuconazole 3.6 Select (equal to 1.13 lbs. of tebuconazole) per acre per year.
- Do not make more than 4 applications per year at highest rate.
- Do not apply to bearing fruit trees or vegetables.

## **LABELED TURF DISEASES INCLUDE**

Anthracnose  
Bermuda Grass Decline  
Brown Patch  
Brown Ring Patch  
Copper Spot  
Corticium Red Thread  
Dollar Spot  
Fusarium Patch  
Gray Leaf Spot  
Gray Snow Mold  
Large Patch  
Microdochium Patch  
Necrotic Ring Spot  
Pink Patch  
Pink Snow Mold  
Powdery Mildew  
Red Thread  
Rhizoctonia Blight  
Rusts  
Spring Dead Spot  
Stripe Smut  
Summer Patch  
Take All Patch  
Typhula Blight  
Zoysia Patch

## **LABELED ORNAMENTAL DISEASES INCLUDE**

Anthracnose  
Black Spot  
Leaf Spot  
Petal Blight  
Powdery Mildew  
Rusts  
Rhizoctonia Stem and Root Rot  
Southern Blight  
Tulip Fire



Prime Source, LLC  
4609 E. Boonville-New Harmony Road  
Evansville, IN 47725  
www.Prime-SourceLLC.com

**Download a  
specimen label at  
Prime-SourceLLC.com**