







We create chemistry



Tower® herbicide

Adaptive preemergent control

Tower herbicide delivers selective preemergent control of annual grasses, sedges and small-seeded broadleaf weeds in golf course turf, lawns and ornamentals.

Challenging Weeds	Adaptive Utility	Premium Tank-Mix Partner	Enhanced Poa Defense
			
Preemergent control of goosegrass, doveweed, sedges, spurge, <i>Poa annua</i> and more	Flexible rate range for warm-season and cool-season turf and short reseeding intervals	Readily mixes with preemergent and postemergent herbicides to expand spectrum of control	Residual control and early postemergent activity on <i>Poa annua</i>


Advantages of Tower herbicide

- Selective preemergent control of annual grassy weeds, broadleaf weeds and sedges
- Inhibits shoot growth to control weeds as they germinate without root pruning effects on desirable turf
- Non-staining, liquid formulation readily mixes with other herbicides to expand spectrum of control
- Reliable activation with just 1/4" irrigation or rain
- Can be impregnated or coated onto dry bulk granular fertilizers

Adaptive residual protection

- Flexible rate range allows for applications to actively-growing warm- and cool-season turfgrass
- Over-the-top tolerance on 100+ ornamental plant species in landscape beds
- Tank-mix compatible with postemergent herbicides for extended residual control on bareground and gravel sites

Goosegrass residual control



Turf best use recommendations

- Apply to dense and well-established turfgrass
- Make applications spring through fall when soil temperatures are above 55°F
- Use 21 fl oz / A on cool-season turf types (bentgrass, Kentucky bluegrass, fine fescue, perennial ryegrass and tall fescue)
- Use 21 to 32 fl oz / A on warm-season turf types (bahiagrass, bermudagrass, buffalograss, centipedegrass, kikuyugrass, St. Augustinegrass, seashore paspalum and zoysiagrass)
- Water-in application to rinse residue from foliage and activate herbicide barrier
- Make sequential applications 5 to 8 weeks apart

See product label for full use instructions

HRAC Group	Active Ingredients (%)	Usage Rate	Formulation	Packaging	Key Weeds Controlled
15	Dimethenamid-P (63.9%)	21 to 32 fl oz	Emulsifiable concentrate (EC)	2 x 2.5 gal	Goosegrass, doveweed, annual sedges, spurge, annual bluegrass, Florida pusley, eclipta, <i>Poa annua</i>

Always read and follow label directions.

betterturf.basf.us

Products may not be registered for sale or use in all states. Please check with your state or local Extension Service. Refer to label for registered uses. It is a violation of Federal law to use a product in a manner inconsistent with its labeling.

Tower is a registered trademark of BASF. Ronstar is a registered trademark of Environmental Science.
©2024 BASF Corporation. All rights reserved. PSS 24-30-423