

- Kills Broadleaf Weeds Like Clover, Dandelion, and Oxalis
- Prevents Crabgrass and Fertilizes

ACTIVE INGREDIENT: 2,4-D, 2-ethylhexyl ester ... Mecoprop-p 0.071% ..0.160% OTHER INGREDIENTS: ... 98.494% TOTAL... .100.000% THIS PRODUCT CONTAINS: 14.70 lb 2,4-dichlorophenoxyacetic acid equivalent per ton or 0.735% (CAS 1928-43-4)

3.34 lb (+)-(R)-2-(2 methyl-4-chlorophenoxy)propionic acid equivalent per ton or 0.167% (CAS 16484-77-8) 1.42 lb 3,6-dichloro-o-anisic acid equivalent per ton or 0.071% (CAS 1918-00-9) 3.20 lb dithiopyr per ton or 0.160% (CAS 97886-45-8) Isomer Specific By AOAC Methods.

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS, FIRST AID, AND DIRECTIONS FOR USE.

New York State: Do not exceed one application at the recommended rate of 3.6 pounds of product per 1,000 sq.ft. per year.

Net Weight 13 LBS. (5.9 KG)





Product Questions? 855-270-4776 available on the internet at: http://www.aapfco.org/metals.htm Information regarding the contents and levels of metals in this product is EPA Reg. No. 2217-913-7401 • EPA Est. No. 7401-TX-1 Visit Us At: www.tertilome.com BONHAM, TEXAS 75418 230 FM 87

ferti•lome® and Weed-Out® are registered trademarks of Voluntary Purchasing Groups, Inc. be varied by any written or verbal statements or agreements at the point of sale or elsewhere. terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these MHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, INCLUDING ANY WARRANTY OR MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES,

waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain. container. If empty: Place in trash or offer for recycling if available. If partly filled: Call your local solid PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this sunlight. Do not store near food feed, seeds, or other pesticides. PESTICIDE STORAGE: Keep in original container in a cool, dry area, away from extreme heat and direct Do not contaminate water, tood, or teed by storage and disposal.

	STORAGE AND DISPOSAL						
	The above spreader settings are approximate. The age and condition of spreader, speed of operation ind evenness of terrain may require slightly different settings for desired coverage.						
Γ	91	15	Þ١	ferti•lome®/Earthway Ev-N-Spred (Broadcast)			
	Þ	3-1/2	3	Scotts/Republic Speedy Green (Broadcast)			
	3-3/4	3-1/4	2-3/4	Scotts®/Republic Accugreen (Drop)			
	SPREADER SETTINGS Lbs. of product per 1,000 sq. ft. 2.5 3.0 3.6.			SPREADER			

and adjust to apply at the rate of 1 pound of product per 400 square feet. • For 2.5 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 400 square feet and adjust to apply at the rate of 1 pound of product per 275 square feet. • For 3.6 pounds of product per 1,000 square feet of turigrass: Check your spreader on 275 square feet CALIBRATION INSTRUCTIONS: 30 days between applications. The maximum seasonal rate is 7.2 pounds of product per 1,000 sq.ft.

application. The maximum number of broadcast applications is limited to 2 per year with a minimum of Other States: The maximum broadcast application rate is 3.6 pounds of product per 1,000 sq.ft. per New York State: Do not exceed one application at the recommended rate of 3.6 pounds of product per Apply 3.0 pounds of product per 1,000 square feet of established turfgrass. For hybrid bermudagrass, bahiagrass and buffalograss: Apply 2.5 to 3.6 pounds of product per 1,000 square feet of established turtgrass. For Kentucky bluegrass, fescue spp., perennial ryegrass, common bermudagrass, and zoysiagrass: poor weed control or turfgrass injury may result. Use a drop or rotary spreader designed to apply granular fertillizers. Avoid skips or excessive overlaps as

HOW MUCH TO USE ornamental shrubbery or desirable plants. Apply this product when the wind is calm. Avoid spreading granules onto flowers, vegetables, and or disturbed for 24 hours after application. Delay mowing for 3 days after application. wet from dew or rain; poor results will occur if it is applied to dry lawns. The lawn should not be watered This product needs to stick to the leaves of broadleaf weeds and should be applied when the grass is

present, young and actively growing under adequate soil moisture and favorable temperatures (50° to 80° F). For broadleaf weed control: Best results are obtained during the spring and fall when broadleaf weeds are this product is applied before the first tiller stage of crabgrass. (smooth, southern, and large) and other susceptible annual grasses. This product may also provide early post emergent control of crabgrasss. However, post emergent crabgrass control will be obtained only when For crabgrass and other listed grasses: This product is a preemergent herbicide to control crabgrass

label extends into early summer, then a second application can be made at 6 to 10 weeks after the initial For the best results in the spring, apply this product one to two (1 to 2) weeks before the expected This product may be applied as a single application or as split applications in the spring, early summer, or fall.

ΥЛЬЬ	ΟT	ΝΞ

E 320	*7.5% Slowly Available Nitrogen from Polymer Coated Sulfur Coated Urea
muibo2 ,dast	Derived From: Ammonium Sulfate, Polymer Coated Sulfur Coated Urea, Muriate of Pol Borate, Copper Sulfate, Iron Sulfate, Manganese Sulfate, Zinc Sulfate.
	0.05% Water Soluble Zinc (Zn)
%90.0	(nS) oniS
	0.05% Water Soluble Manganese (Mn)
%90.0	Manganese (Mn)
	0.10% Water Soluble Iron (Fe)
%01.0	(9∃) no¹l
	0.05% Water Soluble Copper (Cu)
%90.0	Copper (Cu)
%20.0	Boron (B)
% \u03b4	Soluble Potash(K ₂ O)

GUARANTEED ANALYSIS

52%

ade at a 6 to 10 week	egrass, two applications m	f barnyardgrass and goos	er acceptable control o prval are needed
Wild sweet potato Wild sweet potato Woodsorrel, creeping Woodsorrel, yellow (Oxalis) Woolly croton Wornseed Wornseed Wornseed Wornseed Yerrow, common Yerrow, western Yerrow, western	Smartweed, pale Smartweed, pennsylvania Smooth ofalf-flower Smooth ofalf-flower Sorrel, red (sheep) Sowthistle, spiny Sowthistle, spiny Spanish needles Speedwell, common Speedwell, common	Matchweed Mexicanweed Moneywork Monningglory, ivyleaf Morningglory, ivyleaf Morningglory, ived Morningglory, sell Morningglory, sell Morningglory, sell Mustard, ball Mustard, black Mustard, black Mustard, black Mustard, black	ver, white sweet wer, white sweet side weet side wer, white sweet side were side were side were side wer, sping benry eping beggarweed eping beggarweed sile) weed side week side were wer were wer were were were were

Wild strawberry	Smartweed, ladysthumb	Marcela	Clover, strawberry		
Wild rape	Shepardspurse	Mallow, venice	Clover, red		
Wild radish	Scarlet pimpernel	Mallow, dwarf	Clover, rabbit foot		
Wild parsnip	Rough cinquefoil	Mallow, common	Clover, hop		
noino bliW	Redstem filaree	Mallow, bristly	Clover, alyce		
rocket)	Ragweed, western	Mallow, alkali	Clover, crimson		
Wild mustard (yellow	Ragweed, lancleaf	resbedeza	Cinquetoil		
Wild marigold	Hagweed, common	rampadnarters	Chicory		
Wild lettuce	Ragweed, bur	Kochia	Chickweed, sticky		
Wild garlic	Pusley, large flower	Knotweed, prostrate	Chickweed, mouseear		
Wild four-o-clock	Pusley, florida	Knawel	Chickweed, common		
Wild carrot	Pusley, brazil	Johnny-Jumpup violet	Chamber bitter		
Wild buckweat	Purslane, common	Jimsonweed	gsugejion)		
Wild aster	Puncturevine	Horseweed	Catsear, spotted (false		
Western salsify	Prostrate knotweed	Horsenettle	Catnip dintsO		
Western clematis	Prickly sida	Hoary cress	Carpetweed		
Virginia creeper	plant)	Henbit	Carolina geranium (wild)		
Vetch, wild	Prickly lettuce (compass	Hemp	Buttonweed, Virginia		
Vetch, narrowleaf	Prairie sunflower	Heartleaf drymary	(boorloe)		
Vetch, milk	Бокемеед	Healall	Buttonweed, common		
Vetch, hairy	Polson oak	Hawkweed, yellow	Buttercup, creeping		
Vetch, bird	Poison ivy	Hawkweed, orange	Buttercup, bulbous		
Vervain, tall	Plantain, woolly	Hawkweed, mouseear	Burweed, lawn (spurweed)		
Vervain, prostrate	Plantain, slender	Hawkweed, meadow	Burdock		
Vervain, hoary	Plantain, narrowleaf	Hairy fleabane	Burclover		
Vervain, blue	Plantain, hoary	beawmut	Bullnettle		
Velvetleaf	Plantain, buckhorn	Groundsel	Buckhorn		
Irailing crownvetch	Plantain, broadleat	Ground Ivy	Broomweed		
I histle, yellow (horrible)	Plantain, bracted	Goldenrod	Bristly oxtongue		
I histle, wavyleat	Plantain, blackseed	Galinsoga, smallflower	Bristly extensions		
I histle, scotch	Plains coreopsis (tickseed)	Galinsoga, hairy	Brassbuttence		
Thistle, Russian	Pineapple weed	Fleabane, rough	Blood flower milkweed		
Thistle, plumeless	Pigweed, tumble	(annual)	Black medic		
I histle, platte	Pigweed, smooth	Fleabane, daisy	Black-eyed Susan		
Thistle, musk	Pigweed, Russian	Filaree, whitestem	Bitterweed		
Thistle, flodman (prairie)	Pigweed, redroot	Filaree, redstem	Bittercress, hairy		
Thistle, canada	Pigweed, prostrate	Fiddleneck	Bitter wintercress		
Thistle, bull	(Peppergrass)	False sunflower	Bindweed, Hedge		
I histle, blessed	Pepperweed, Virginia	Falsetlax	Bindweed, Field		
lanweed	Pepperweed, perennial	Elderberry	Betony, Florida		
Iansy ragwort	Pennycress (frenchweed)	Dove weed	Beggarticks, hairy		
Strawberry, India mock	Pearlwort (fronchwood)	(Pennywort)	Begasticks, pries		
Star of Bethlehem	Parsnip	Dollarweed	Bedstraw, Northern		
Spurweed	Parsiey-piert	Dogrennel	Bedstraw, smooth		
Spurge, prostrate	Oxalis	Dogbane	Bedstraw, catchweed		
Spurge, spotted	Nettle, wood	Dock, curly	Austrian fieldcress		
Spiny amaranth	Nettle, tall	Dock, broadleat	Artichoke Austrian fieldcress		
Speedwell, thymeleaf	Nettle, stinging (burning)	Dock broadloaf	Aster, white prairie		
Speedwell, thympolost	Mustard, wormseed	Deadnettle	Aster, purple		
Speedwell, slender	Mustard, white	Description In the control of the co	Aster, white heath		
Speedwell, purslane	Mustard, tumble	Dandelion, talse	Aster, heath		
Speedwell, persian	Mustard, tansy	Dandelion, common	Amaranth, slender Aster, heath		
Speedwell, ivyleaf	Mustard, indian	Daisy, oxeye	Amaranth, purple		
tealyyi llawbaaq2	neibai baetauM	evevo vaisf	Amaranth purple		
		odujedi weeus.	in nansilabisa io gimen M		
A listing of established broadleaf weeds:					
	yellow)		,		
(crowfootgrass)	Foxtail (green and	Barnyardgrass*	(Poa annua)		
202162222	200160010	0001606110011111	00016001010011111		

Annual ryegrass Annual bluegrass* Annual Grass Weeds When applied as directed, this product will control and/or suppress* the following weeds: MEEDS CONTROLLED

• Do not use clippings from treated areas for composting or for mulching vegetables or fruit trees. Do not exceed specified dosage for any area. area(s) to the point of runoff or apply when raining or when rain is expected that day. gutter, or street back onto the treated area. To prevent product run-off, do not over water the treated • Apply this product directly to your lawn, and sweep any product landing on the driveway, sidewalk, Do not apply when windy. • Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches.

desirable trees and shrubs, since injury may result. • Do not use this product on or near desirable plants, including within the drip line of the roots of OTHER USE PRECAUTIONS: weakened by these stresses to recover before application of this product. been weakened by stress due to mechanical operations, chemical injury, disease, insects, pests, excessive temperatures, or lack of moisture may increase the chances for turigrass injury. Allow turigrass The use of this product on lawns and ornamental turfgrasses that are not well established, or have GUIDELINES TO FOLLOW: Augustinegrass in Florida. Do not use this product on turfgrass where desirable clovers are present.

Do not use this product on dichondra, carpetgrass, or colonial bentgrass. Do not use this product on St. Zoysiagrass Buffalograss Fine Fescues Bermudagrass Perennial Ryegrass Rahiagrass Kentucky Bluegrass ESTABLISHED COOL SEASON TURFGRASS ESTABLISHED WARM SEASON TURFGRASS This product may be used on the following established turt species:

WHERE TO USE

thistle, and clover. A partial list of broadleaf weeds is presented on this label.

• The post emergent broadleaf herbicide will control over 250 lawn weeds including dandelion, chickweed, The fertilizer contains slow release nutrients that may extend the availability for turfgrass uptake. foxtails, and other undesirable annual grasses. • The preemergent herbicide can be applied to established turigrass to control crabgrass, goosegrass, This product contains three components: a preemergent herbicide, a lawn fertilizer, and a post emergent This product is formulated to provide control of most annual grasses and certain broadleaf weeds

• Not for use on turt being grown for sale or other commercial use as sod, or for commercial seed or for treated area until dusts have settled. Keep people and pets out of the area during application. Do not allow people or pets to enter the Do not apply this product in a way that will contact any person or pet, either directly or through drift. GENERAL PRECAUTIONS AND RESTRICTIONS: It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DIBECTIONS FOR USE

result in groundwater contamination. Application around a cistern or well may result in contamination of use or rnis chemical in areas where soils are permeable, particularly where the water table is shallow, may This chemical has properties and characteristics associated with chemicals detected in groundwater. The apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate. This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not ENVIRONMENTAL HAZARDS for treatment. For additional information in case of emergency, call toll free 1-877-800-5556. Have the product container or label with you when calling a poison control center or doctor or going • Call a poison control center or doctor for treatment advice. Rinse skin immediately with plenty of water for 15 - 20 minutes. Take off contaminated clothing. If On Skin Or

• Do not give anything by mouth to an unconscious person. • Do not induce vomiting unless told to do so by a poison control center or Have person sip a glass of water if able to swallow. • Call a poison control center or doctor immediately for treatment advice. :bewellowed: • Call a poison control center or doctor for treatment advice. Hemove contact lenses, it present, after the first 5 minutes, then continue • Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. If In Eyes:

> HAZARDS TO HUMANS AND DOMESTIC ANIMALS STOP! DIRECTIONS CAREFULLY.
>
> DIRECTIONS CAREFULLY.

tobacco. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using

Weed-Out® Pro Turf

24% Urea Nitrogen*

Total Nitrogen (N)..

1% Ammoniacal Nitrogen

KEEP OUT OF REACH OF CHILDREN