

APPROVED

By Kyndra Lemke at 12:54 pm, Jan 23, 2009

Weed-Out Broadleaf Weed Control 10925



Weed-Out

BROADLEAF WEED CONTROL
CONTAINS TRIMEC® HERBICIDE

- Controls: Dandelion, Chickweed, Knotweed, Plantain, Henbit, Spurge And Many Other Broadleaf Weeds.
- For Use On Bahia, Bermuda, Centipede, Zoysia, Kentucky Bluegrass, Perennial Rye, Fescue And Bentgrass Lawns.



ACTIVE INGREDIENTS:

Dimethylamine salt of 2,4-dichlorophenoxyacetic acid.....	1.65%
Dimethylamine salt of (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid.....	0.40%
Dimethylamine salt of dicamba (3,6-dichloro-o-anisic acid).....	0.18%
INERT INGREDIENTS:	97.77%
TOTAL	100.00%

THIS PRODUCT CONTAINS:

27.4 lbs. 2,4-dichlorophenoxyacetic acid equivalent per ton or 1.37%.
 6.6 lbs. (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per ton or 0.33%.
 3.0 lbs. 3,6-dichloro-o-anisic acid equivalent per ton or 0.15%.
 Isomer Specific by AOAC Methods.
 Contains the single Isomer form of Mecoprop-p.
 TRIMEC® is a registered trademark of PBI/Gordon Corporation.

**KEEP OUT OF REACH
OF CHILDREN
CAUTION**

See Back Panel For Additional
Precautionary Statements.

NET WEIGHT 10 LBS. (4.54 KG)

10925

Weed-Out Broadleaf Weed Control



Weed-Out Broadleaf Weed Control



10925

Bottom of Bag

Bottom of Bag



Weed-Out

BROADLEAF WEED CONTROL

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Causes moderate eye irritation. Avoid breathing dust. Avoid contact with skin, eyes or clothing. When using this product, wear long-sleeved shirt, long pants, socks and shoes. After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water.

FIRST AID

If In Eyes:	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a Poison Control Center or doctor for treatment advice.
If On Skin Or Clothing:	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. Call a Poison Control Center or doctor for treatment advice.
If Swallowed:	<ul style="list-style-type: none"> Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a Poison Control Center or doctor. Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a Poison Control Center or doctor or going for treatment. You may also call toll-free, 1-877-800-5556 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment area until dust has settled.

STORAGE AND DISPOSAL

STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals. Do not store near food or feedstuffs. **PESTICIDE DISPOSAL:** Securely wrap bag in several layers of newspaper and discard in trash. **DISPOSAL: If Empty:** Do not reuse bag. Place in trash or offer for recycling if available. **If Partly Filled:** Call your Local Solid Waste Agency for disposal instructions. Never place unused product down any indoor or out door drain.

USE PRECAUTIONS

Do not apply this product to St. Augustinegrass, Carpetgrass, Dichondra, Ornamentals, or Lawns containing desirable Clovers. Avoid spreading granules onto Landscape, Plantings, Flowers, Vegetables or Ornamentals. Do not spread over exposed roots of Trees and Ornamentals. Do not exceed specified dosages for any area; be particularly careful within drip-line of Tree and other Ornamental Species. Treated areas may be seeded 3 to 4 weeks after application. Application of this product to newly seeded areas should be delayed until the third mowing.

WHERE TO USE

ferti·lome® WEED-OUT® BROADLEAF WEED CONTROL is a **Trimec®** complex, a combination of three proven Weed Killers – 2, 4-D, Mecoprop and Dicamba. Together, they control a wide range of Lawn Weeds, including Dandelion, Chickweed, Knotweed, Plantains, Henbit, Spurge and many others found in Home Lawns. Established turf of Kentucky Bluegrass, Bentgrass, Bahiagrass, Centipedegrass, Fescues, Bermudagrass, Zoysiagrass, and Perennial Ryegrass may be treated.

HOW MUCH TO APPLY

For Kentucky Bluegrass, Bentgrass, Bahiagrass, Centipedegrass, Fescues, Perennial Ryegrass, Bermudagrass, and Zoysiagrass: Apply 2.0 pounds of product per 1,000 square feet of turfgrass. A second or follow-up application may be required for dense infestations of Weeds or for adverse seasonal conditions. The second application of 2.0 pounds of product per 1,000 square feet should be used 30 days after the first application.

The maximum application rate to turf is 1.2 pounds 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year.

- Mow lawn to normal height 1 to 2 days before application.
- Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering (see step 4).
- Apply when Weeds are young and actively growing, preferably in the morning when dew is on the Grass. If Grass is not moist at time of application, sprinkle lightly with water to hold the material and prevent dust drift onto nontarget Plants.
- Do not wash from Weed leaves for 1 to 2 days after application. At this time a thorough watering should be made.

CALIBRATION INSTRUCTIONS

A uniform distribution of product on the treated area is essential. Choose a calm day to ensure uniform coverage and to avoid spreading granules onto Flowers, Vegetables, and Ornamental Shrubbery. Do not exceed the application rate for any area and use care in the application of this product within the dripline of Trees, Shrubs, and other Ornamentals. Drop spreaders and rotary spreaders should be calibrated for uniform applications of this product.

For 2.0 lbs. of product per 1,000 sq. ft. of turfgrass: Check your spreader on 500 sq. ft. and adjust to apply at the rate of 1 lb. of product per 500 sq. ft.

SPREADER GUIDE ONE BAG WILL COVER UP TO 5,000 SQUARE FEET

SPREADER	SPREADER SETTINGS
ferti·lome® Plus Iron Broadcast/Earthway 2850	13
ferti·lome® Plus Iron Prof. Broadcast/Earthway 2800A	13
ferti·lome® Deluxe Broadcast/Earthway 2050 S.U.	13
ferti·lome® By SPYKER	4.4
Scott Broadcast/Speedy Green	7
Republic/Scotts Broadcast	6
Republic Drop	7
Precision Broadcast/Red Devil	8
Precision Drop FH-22D	7



These settings can be used only as a guide, they are not absolute. The speed at which the applicator walks, spreader condition and the texture of the turf will alter the rate of application. If product remains in spreader after application to recommended square foot coverage, spread remainder over same area going at right angles (See Diagram).

THE FOLLOWING BROADLEAF WEEDS ARE CONTROLLED BY THIS PRODUCT:

Annual Yellow Sweetclover	Burclover	Daisy, Fleabane	Knawel	Poison Ivy	Sorrel	White Mustard
Aster	Burdock	Daisy, Oxeye	Knotweed	Poison Oak	Sowthistle	Wild Aster
Austrian Fieldcress	Burning Nettle	Dandelion	Kochia	Pokeweed	Spanishneedles	Wild Buckwheat
Bedstraw	Burweed	Dichondra	Lambsquarters	Poorjoe	Speedwell	Wild Carrot
Beggarticks	Buttercup	Dogfennel	Lespedeza	Prairie Sunflower	Spiny Amaranth	Wild Four-O'Clock
Betony, Florida	Buttonweed	Dollarweed	Mallow	Prickly Lettuce	Spiny Cocklebur	Wild Garlic
Bindweed, Field	Canada Thistle	False Dandelion	Matchweed	Prickly Sida	Spiny Sowthistle	Wild Geranium
Bird Vetch	Carolina Geranium	Falseflax	Mexicanweed	Prostrate Knotweed	Spotted Catsear	Wild Lettuce
Bitter Wintercress	Carpetweed	False Sunflower	Milk Vetch	Prostrate Pigweed	Spotted Spurge	Wild Marigold
Bittercress, Hairy	Catchweed Bedstraw	Fiddleneck	Morningglory	Prostrate Spurge	Spurweed	Wild Mustard
Bitterweed	Catnip	Florida Pusley	Mouseear Hawkweed	Prostrate Vervain	Stinging Nettle	Wild Onion
Black-eyed Susan	Catsear	Frenchweed	Mugwort	Puncturevine	Strawberry, India Mock	Wild Parsnip
Black Medic	Chickweed, Common	Galinsoga	Musk Thistle	Purslane, Common	Tall Nettle	Wild Radish
Black Mustard	Chickweed, Mouseear	Goldenrod	Mustard	Ragweed	Tall Vervain	Wild Rape
Blackseed Plantain	Chicory	Ground Ivy	Narrowleaf Plantain	Red Sorrel	Tansy Ragwort	Wild Strawberry
Blessed Thistle	Cinquefoil	Gumweed	Narrowleaf Vetch	Redroot Pigweed	Tansy Mustard	Wild Sweet Potato
Bloodflower Milkweed	Clover, Crimson	Hairy Fleabane	Nettle	Redstem Filaree	Tanweed	Wild Vetch
Blue Lettuce	Clover, Hop	Hawkweed	Orange Hawkweed	Rough Cinquefoil	Thistle	Woodsorrel
Blue Vervain	Clover, Red	Healall	Oriental Cocklebur	Rough Fleabane	Trailing Crownvetch	Woolly Croton
Bracted Plantain	Clover, Strawberry	Heartleaf Drymary	Oxalis	Russian Pigweed	Tumble Mustard	Woolly Morningglory
Bristly Oxtongue	Clover, Sweet	Heath Aster	Parsley-piert	Russian Thistle	Tumble Pigweed	Woolly Plantain
Broadleaf Dock	Clover, White	Hedge Bindweed	Parsnip	Scarlet Pimpernel	Velvetleaf	Wormseed
Broadleaf Plantain	Cockle	Hedge Mustard	Pearwort	Scotch Thistle	Venice Mallow	Yarrow
Broomweed	Cocklebur	Henbit	Pennycress	Sheep Sorrel	Virginia Buttonweed	Yellow Rocket
Buckhorn	Common Mullein	Hoary Cress	Pepperweed	Sheepspurge	Virginia Creeper	Yellowflower Pepperweed
Buckhorn Plantain	Creeeping Jenny	Hoary Plantain	Pigweed	Slender Plantain	Virginia Pepperweed	and other Broadleaf Weeds
Bulbous Buttercup	Cudweed	Hoary Vervain	Pineywoods Bedstraw	Smallflower Galinsoga	Wavyleaf Bullthistle	
Bull Thistle	Curly Dock	Horsenettle	Plains Coreopsis	Smooth Dock	Western Clematis	
Bullnettle	Daisy, English	Jimsonweed	Plantain	Smooth Pigweed	Western Salsify	

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment will vary, Seller makes no claims other than those stated on this label.

Distributed By:
VOLUNTARY PURCHASING GROUPS, INC.
230 FM 87
BONHAM, TEXAS 75418
EPA Reg. No. 2217-657-7401 • EPA Est. No. 7401-TX-1
Visit Us At: www.fertilome.com



1-09

Bottom of Bag

Bottom of Bag



**Weed-Out Broadleaf
Weed Control**



10.875 "x 22.5"