

# ferti·lome<sup>®</sup>

**Kills Weeds &  
Prevents Crabgrass While  
Feeding Your Lawn**

# Weed Out<sup>®</sup> Pro Turf

## 25-0-4

- Kills Broadleaf Weeds Like Clover, Dandelion, and Oxalis
- Prevents Crabgrass and Fertilizes

**ACTIVE INGREDIENT:**

2,4-D, 2-ethylhexyl ester .....	1.108%
Mecoprop-p .....	0.167%
Dicamba .....	0.071%
Dithiopyr .....	0.160%

OTHER INGREDIENTS: ..... 98.494%  
TOTAL..... 100.000%

**THIS PRODUCT CONTAINS:**

14.70 lb 2,4-dichlorophenoxyacetic acid equivalent per ton or 0.735% (CAS 1928-43-4)  
3.34 lb (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid equivalent per ton or 0.167% (CAS 16484-77-8)  
1.42 lb 3,6-dichloro-o-anisic acid equivalent per ton or 0.071% (CAS 1918-00-9)  
3.20 lb dithiopyr per ton or 0.160% (CAS 97886-45-8)  
Isomer Specific By AOAC Methods.

**KEEP OUT OF REACH OF CHILDREN**

### CAUTION

**SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY  
STATEMENTS, FIRST AID, AND DIRECTIONS FOR USE.**

New York State: Do not exceed one application at the recommended rate of  
3.6 pounds of product per 1,000 sq. ft. per year.

**Net Weight 13 LBS. (5.9 KG)**

**Weed Out<sup>®</sup> Pro Turf**

**Weed Out<sup>®</sup> Pro Turf**

**Covers Up To  
5,000 Square Feet**

# Weed Out<sup>®</sup> Pro Turf



KEEP OUT OF REACH OF CHILDREN

# Weed Out® Pro Turf

**STOP!** READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** Causes moderate eye irritation. Avoid contact with eyes or clothing. Wear protective eyewear. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

## FIRST AID

If In Eyes:	<ul style="list-style-type: none"> <li>Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.</li> <li>Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
If Swallowed:	<ul style="list-style-type: none"> <li>Call a poison control center or doctor immediately for treatment advice.</li> <li>Have person sip a glass of water if able to swallow.</li> <li>Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>Do not give anything by mouth to an unconscious person.</li> </ul>
If On Skin Or Clothing:	<ul style="list-style-type: none"> <li>Take off contaminated clothing.</li> <li>Rinse skin immediately with plenty of water for 15 - 20 minutes.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>

Have the product container or label with you when calling a poison control center or doctor or going for treatment. For additional information in case of emergency, call toll free 1-877-800-5556.

## ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

### GENERAL PRECAUTIONS AND RESTRICTIONS:

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.
- Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed or for research purposes.

## PRODUCT DESCRIPTION

This product is formulated to provide control of most annual grasses and certain broadleaf weeds commonly found in home lawns.

This product contains three components: a preemergent herbicide, a lawn fertilizer, and a post emergent broadleaf weed killer.

- The preemergent herbicide can be applied to established turfgrass to control crabgrass, goosegrass, foxtails, and other undesirable annual grasses.
- The fertilizer contains slow release nutrients that may extend the availability for turfgrass uptake.
- The post emergent broadleaf herbicide will control over 250 lawn weeds including dandelion, chickweed, thistle, and clover. A partial list of broadleaf weeds is presented on this label.

## WHERE TO USE

This product may be used on the following established turf species:

ESTABLISHED COOL SEASON TURFGRASS	ESTABLISHED WARM SEASON TURFGRASS
Kentucky Bluegrass	Bahiagrass
Perennial Ryegrass	Bermudagrass
Fine Fescues	Buffalograss
Tall Fescue	Zoysiagrass

Do not use this product on dichondra, carpetgrass, or colonial bentgrass. Do not use this product on St. Augustinegrass in Florida. Do not use this product on turfgrass where desirable clovers are present.

### GUIDELINES TO FOLLOW:

The use of this product on lawns and ornamental turfgrasses that are not well established, or have been weakened by stress due to mechanical operations, chemical injury, disease, insects, pests, excessive temperatures, or lack of moisture may increase the chances for turfgrass injury. Allow turfgrass weakened by these stresses to recover before application of this product.

### OTHER USE PRECAUTIONS:

- Do not use this product on or near desirable plants, including within the drip line of the roots of desirable trees and shrubs, since injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches.
- Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans.
- Do not apply when windy.
- Apply this product directly to your lawn, and sweep any product landing on the driveway, sidewalk, gutter, or street back onto the treated area. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day.
- Do not exceed specified dosage for any area.
- Do not use clippings from treated areas for composting or for mulching vegetables or fruit trees.

## WEEDS CONTROLLED

When applied as directed, this product will control and/or suppress\* the following weeds:

Annual Grass Weeds			
Annual bluegrass* (Poa annua)	Annual ryegrass Barnyardgrass*	Crabgrass Foxtail (green and yellow)	Goosegrass* (crowfootgrass)

A listing of established broadleaf weeds:

Amaranth, purple	Daisy, oxeye	Mustard, indian	Speedwell, ivyleaf
Amaranth, slender	Dandelion, common	Mustard, tansy	Speedwell, persian
Aster, heath	Dandelion, false	Mustard, tumble	Speedwell, purslane
Aster, white heath	Day flower	Mustard, white	Speedwell, slender
Aster, purple	Deadnettle	Mustard, wormseed	Speedwell, snow
Aster, white prairie	Dichondra	Nettle, stinging (burning)	Speedwell, thymeleaf
Artichoke	Dock, broadleaf	Nettle, tall	Spiny amaranth
Austrian fieldcress	Dock, curly	Nettle, wood	Spurge, spotted
Bedstraw, catchweed	Dogbane	Oxalis	Spurge, prostrate
Bedstraw, smooth	Dogfennel	Parsley-piert	Spurweed
Bedstraw, Northern	Dollarweed	Parsnip	Star of Bethlehem
Bedstraw, pinewoods	(Pennywort)	Pearlwort	Strawberry, India mock
Beggarticks, hairy	Dove weed	Pennycress (frenchweed)	Tansy ragwort
Betony, Florida	Elderberry	Pepperweed, perennial	Tanweed
Bindweed, Field	Falseflax	Pepperweed, Virginia	Thistle, blessed
Bindweed, Hedge	False sunflower	(Peppergrass)	Thistle, bull
Bitter wintercress	Fiddleneck	Pigweed, prostrate	Thistle, canada
Bittercress, hairy	Filaree, redstem	Pigweed, redroot	Thistle, fiodman (prairie)
Bitterweed	Filaree, whitemstem	Pigweed, Russian	Thistle, musk
Black-eyed Susan	Flabane, daisy	Pigweed, smooth	Thistle, platte
Black medic	(annual)	Pigweed, tumble	Thistle, plumeless
Blood flower milkweed	Flabane, rough	Pineapple weed	Thistle, Russian
Blue lettuce	Galinsoga, hairy	Plains coreopsis (tickseed)	Thistle, scotch
Brassbuttons	Galinsoga, smallflower	Plantain, blackseed	Thistle, wavyleaf
Bristly oxtongue	Goldenrod	Plantain, bracted	Thistle, yellow (horrible)
Broomweed	Ground ivy	Plantain, broadleaf	Trailing crownwetch
Buckhorn	Groundsel	Plantain, buckhorn	Velvetleaf
Bullnettle	Gumweed	Plantain, hoary	Vervain, blue
Burclover	Hairy fleabane	Plantain, narrowleaf	Vervain, hoary
Burdock	Hawkweed, meadow	Plantain, slender	Vervain, prostrate
Burnweed, lawn (spurweed)	Hawkweed, mouseear	Plantain, woolly	Vervain, tall
Buttercup, bulbous	Hawkweed, orange	Poison ivy	Vetch, bird
Buttercup, creeping	Hawkweed, yellow	Poison oak	Vetch, hairy
Buttonweed, common (poorjoe)	Healall	Pokeweed	Vetch, milk
Buttonweed, Virginia	Heartleaf drymary	Prairie sunflower	Vetch, narrowleaf
Carolina geranium (wild)	Hemp	Prickly lettuce (compass plant)	Vetch, wild
Carpetweed	Henbit	Prickly sida	Virginia creeper
Catnip	Hoary cress	Prostrate knotweed	Western clematis
Catsear, spotted (false dandelion)	Horse-nettle	Puncturevine	Western salsify
Chamber bitter	Horseweed	Purslane, common	Wild aster
Chickweed, common	Johnny-jumpup violet	Pusley, brazil	Wild buckwheat
Chickweed, mouseear	Knawel	Pusley, florida	Wild carrot
Chickweed, sticky	Knotweed, prostrate	Pusley, large flower	Wild four-o'clock
Chicory	Kochia	Ragweed, bur	Wild garlic
Cinquefoil	Lambsquarters	Ragweed, common	Wild lettuce
Clover, crimson	Lespedeza	Ragweed, lanceleaf	Wild marigold
Clover, alyce	Mallow, alkali	Ragweed, western	Wild mustard (yellow rocket)
Clover, hop	Mallow, bristly	Redstem filaree	Wild onion
Clover, rabbit foot	Mallow, common	Rough cinquefoil	Wild pansip
Clover, red	Mallow, dwarf	Scarlet pimpernel	Wild radish
Clover, strawberry	Mallow, venice	Shepardspurse	Wild rape
	Marcela	Smartweed, ladythumb	Wild strawberry

Clover, white	Matchweed	Smartweed, pale	Wild sweet potato
Clover, white sweet	Mexicanweed	Smartweed, pennsylvania	Wild violet
Clover, yellow sweet	Moneywort	Smooth chaff-flower	Woodsorrel, creeping
Cockle	Morningglory, bigroot	Smoothdock	Woodsorrel, violet
Cocklebur	Morningglory, ivyleaf	Sorrel, red (sheep)	Woodsorrel, yellow (Oxalis)
Cocklebur, oriental	Morningglory, mexican	Sowthistle, annual	Woolly croton
Cocklebur, spiny	Morningglory, red	Sowthistle, spiny	Wormseed
Common mullein	Morningglory, tall	Spanish needles	Yarrow, common
Corn spurry	Morningglory, wolly	Speedwell, bi-lobed	Yarrow, western
Creeping beggarweed	Mugwort	Speedwell, common	Yarrow, yellow
Creeping Jenny (creeping charlie)	Mustard, ball	Speedwell, corn	
Cudweed	Mustard, black		
Cupid shaving brush	Mustard, blue		
Daisy, English	Mustard, hedge		

\* For acceptable control of Barnyardgrass and Goosegrass, two applications made at a 6 to 10 week interval are needed

# 25-0-4

## GUARANTEED ANALYSIS

Total Nitrogen (N).....	25%
1% Ammoniacal Nitrogen	
24% Urea Nitrogen*	
Soluble Potash(K <sub>2</sub> O).....	4%
Boron (B).....	0.02%
Copper (Cu).....	0.05%
0.05% Water Soluble Copper (Cu)	
Iron (Fe).....	0.10%
0.10% Water Soluble Iron (Fe)	
Manganese (Mn).....	0.05%
0.05% Water Soluble Manganese (Mn)	
Zinc (Zn).....	0.05%
0.05% Water Soluble Zinc (Zn)	

Derived From: Ammonium Sulfate, Polymer Coated Sulfur Coated Urea, Muriate of Potash, Sodium Borate, Copper Sulfate, Iron Sulfate, Manganese Sulfate, Zinc Sulfate.

\*7.5% Slowly Available Nitrogen from Polymer Coated Sulfur Coated Urea F 370

## WHEN TO APPLY

### Established lawns:

This product may be applied as a single application or as split applications in the spring, early summer, or fall.

For the best results in the spring, apply this product one to two (1 to 2) weeks before the expected germination of crabgrass. When the period of germination for crabgrass and other weeds listed on this label extends into early summer, then a second application can be made at 6 to 10 weeks after the initial application.

**For Crabgrass and other listed grasses:** This product is a preemergent herbicide to control Crabgrass (smooth, southern, and large) and other susceptible annual grasses. This product may also provide early post emergent control of Crabgrass. However, post emergent Crabgrass control will be obtained only when this product is applied before the first tiller stage of Crabgrass.

**For broadleaf weed control:** Best results are obtained during the spring and fall when broadleaf weeds are present, young and actively growing under adequate soil moisture and favorable temperatures (50° to 80°F).

## HOW TO APPLY

This product needs to stick to the leaves of broadleaf weeds and should be applied when the grass is wet from dew or rain; poor results will occur if it is applied to dry lawns. The lawn should not be watered or disturbed for 24 hours after application. Delay mowing for 3 days after application.

Apply this product when the wind is calm. Avoid spreading granules onto flowers, vegetables, and ornamental shrubbery or desirable plants.

## HOW MUCH TO USE

Use a drop or rotary spreader designed to apply granular fertilizers. Avoid skips or excessive overlaps as poor weed control or turfgrass injury may result.

### For Kentucky Bluegrass, Fescue spp., Perennial Ryegrass, Common Bermudagrass, and Zoysiagrass:

Apply 2.5 to 3.6 pounds of product per 1,000 square feet of established turfgrass.

### For hybrid Bermudagrass, Bahiagrass and Buffalograss:

Apply 3.0 pounds of product per 1,000 square feet of established turfgrass.

**New York State:** Do not exceed one application at the recommended rate of 3.6 pounds of product per 1,000 sq.ft. per year.

**Other States:** The maximum broadcast application rate is 3.6 pounds of product per 1,000 sq.ft. per application. The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 7.2 pounds of product per 1,000 sq.ft.

## CALIBRATION INSTRUCTIONS:

- For 3.6 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 275 square feet and adjust to apply at the rate of 1 pound of product per 275 square feet.
- For 2.5 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 400 square feet and adjust to apply at the rate of 1 pound of product per 400 square feet.

## SUGGESTED SPREADER SETTINGS

SPREADER	SPREADER SETTINGS Lbs. of product per 1,000 sq. ft.		
	2.5	3.0	3.6
Scotts®/Republic Accugreen (Drop)	2-3/4	3-1/4	3-3/4
Scotts/Republic Speedy Green (Broadcast)	3	3-1/2	4
ferti•lome®/Earthway Ev-N-Spred (Broadcast)	14	15	16

The above spreader settings are approximate. The age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

**PESTICIDE STORAGE:** Keep in original container in a cool, dry area, away from extreme heat and direct sunlight. Do not store near food feed, seeds, or other pesticides.

**PESTICIDE DISPOSAL AND CONTAINER HANDLING:** Nonrefillable container. Do not reuse or refill this container. If empty: Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

## LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OR MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere. ferti•lome® and Weed Out® are registered trademarks of Voluntary Purchasing Groups, Inc.



230 FM 87  
BONHAM, TEXAS 75418  
Visit Us At: www.fertilome.com

EPA Reg. No. 2217-913-7401 • EPA Est. No. 7401-TX-1  
Information regarding the contents and levels of metals in this product is available on the internet at: <http://www.aapfc.org/metals.htm>

Product Questions? 855-270-4776

