



DEPEND ON WEEDMASTER®.

For years, growers and ranchers have counted on WeedMaster® for consistent weed control in sugarcane, pastures, small grains and fallowland. WeedMaster is a selective, post-emerge herbicide that provides tough, long-lasting control of a broad spectrum of annual, biennial and perennial broadleaf weeds. The growth regulating effects of 2,4-D and dicamba interfere with the normal development of weeds, ultimately causing plant death.

COMPETITIVE ADVANTAGES

CONSISTENT CONTROL IN A CONCENTRATED FORMULA

- Achieve outstanding control of brush and broadleaf weeds in a concentrated formula
- Controls tough weeds such as horsenettle, thistles and blackberry
- Safe for desirable grasses
- Pre-emergence control with residual activity

AT-A-GLANCE INFO

EPA REG. NO.	71368-34
ACTIVE INGR.	DMA dicamba salt (12.4%), 2,4-D amine salt (35.7%)
FORMULATION	liquid
CHEM. FAMILY	benzoic acid, phenoxycarboxylic acid
HRAC NO.	4
MODE OF ACT	auxin growth regulator
REI	48 hours
SIGNAL WORD	danger
RESTRICTED	no
PACKAGE SIZE	1 gal container (5 per case), 2.5 gal container (2 per case), 30 gal drum, 265 gal tote, bulk

MORE PRODUCTS EXCLUSIVELY FROM NUFARM

BurnMaster™ | **Scorch™** | **Spitfire™**

 **Nufarm**
Grow a better tomorrow



DEPEND ON WEEDMASTER®.

KEY USES

Conservation Reserve Program (CRP) land
Fallowland
General farmstead
Sorghum
Grass (hay or silage)
Pasture
Rangeland
Sugarcane
Wheat

KEY WEEDS CONTROLLED

(See product label for complete list)

ANNUALS	BIENNIALS & PERENNIALS
Buckwheat, wild	Bindweed, field
Burdock	Bittercress
Buttercup	Chicory
Cockle, cow	Dandelion, common
Cocklebur, common	Dock, curly
Dogfennel	Goldenrod, Missouri
Evening primrose	Groundsel, Texas
Fleabane, annual	Honeysuckle, hairy
Flixweed	Horsenettle, Carolina
Knotweed	Ivy, poison
Kochia	Marshelder
Lambsquarters, common	Mesquite
Marestail (horseweed)	Persimmon, eastern
Morningglory; ivyleaf, tall	Prickly lettuce
Mustards; annual, tansy	Smartweed
Pigweed; prostrate, redroot, smooth, tumble	Sorrel, red
Purslane, common	Thistle, bull
Ragweed; common, lanceleaf, western	Vetch, hairy
Sowthistle	
Shepherd's-purse	
Smartweed, Pennsylvania	
Sneezeweed, bitter	
Thistle, Russian	
Velvetleaf	

USE SUMMARY

RATES: 0.5 pts per acre to 4 to 5 pts/A for control of annual, biennial and perennial weeds. See label for per acre rate according to weed growth stage.

Apply to actively growing weeds as aerial, broadcast, band, or spot spray applications.

Apply with water or sprayable fluid fertilizer as a carrier. Sprayable fluid fertilizer may be used as the carrier in preplant or pre-emergence uses. Post-emergence uses with sprayable fluid fertilizer may be made only on pasture, hayland, or wheat crops.

Surfactants and/or low use rate liquid fertilizers (28-0-0, 32-0-0) or crop oil concentrate may be used with this product to improve burndown of emerged weeds. (Do not use crop oil concentrate for feed crops – refer to crop oil concentrate label for complete directions.)

AERIAL APPLICATION: Use 3 to 10 gals of water/A.

GROUND APPLICATION (BROADCAST): Use 5 to 40 gals per broadcast acre.

See label for complete directions for use.



nufarmpoint.com

For specific application rates, directions, mixing instructions and precautions, read the product label. Please visit www.nufarm.com/us to download a full label.
©2018 Nufarm. Important: Always read and follow label instructions. BurnMaster™, Scorch™, Spitfire® and WeedMaster® are trademarks of Nufarm.



15-AG-0073-D