

This information is for promotional purposes only. Space considerations may require information to be omitted.
Always refer to the actual package label for complete use directions. This product may not yet be available or approved for sale or use in your area.

ZoGuard[™] plus FOR CATS

**READ ENTIRE LABEL BEFORE EACH USE.
USE ONLY ON CATS AND PUPPIES OVER 8 WEEKS OF AGE.**

ZoGuard[™] Plus for Cats contains Fipronil and S-Methoprene (insect growth regulator) to effectively kill fleas (all stages) and ticks (including those that may transmit Lyme disease). The active ingredients in ZoGuard Plus for Cats feature a sustained release formula developed for fast-acting and long-lasting control of fleas, ticks, and chewing lice for 1 month on cats only.

ACTIVE INGREDIENTS:

Fipronil:	9.80%
(S)-methoprene:	11.80%
OTHER INGREDIENTS:	78.40%
TOTAL:	100.0%

KEEP OUT OF REACH OF CHILDREN CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS. CAUTION. Harmful if swallowed. Causes eye irritation. Avoid contact with skin, eyes or clothing. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

HAZARDS TO DOMESTIC ANIMALS. For external use on cats only. Individual sensitivities, while rare, may occur after using any pesticide product on cats. Certain medications can interact with pesticides. Do not treat your cat with more than one pesticide product at a time. Over dosing your cat can result in serious illness and even death. Consult a veterinarian before using on medicated, debilitated, aged, pregnant or nursing cats. Consult a veterinarian before using on cats with known organ dysfunction. If your cat is exhibiting signs of and/or is being treated for skin dermatitis, talk to your veterinarian before applying any topical flea and tick control product.

FIRST AID

If swallowed:	Call a poison control center or a doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
If in eyes:	Flush eyes with plenty of water. Call a doctor if irritation persists.
If on skin:	Wash with plenty of soap and water. Get medical attention if irritation persists.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-844-685-9173 for emergency medical treatment information.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT ALLOW CHILDREN TO APPLY PRODUCT. TO PREVENT HARM TO YOU AND YOUR Cat, READ ENTIRE LABEL BEFORE EACH USE. FOLLOW ALL DIRECTIONS AND PRECAUTIONARY STATEMENTS CAREFULLY. Do not use on cats, rabbits or animals other than cats. Do not allow your cat to ingest this product. Do not use on kittens under 8 weeks of age. Do not apply to cats weighing less than 1.5 lbs. Use entire contents of tube vial on each cat. Do not use multiple tubes on one cat. Weigh your cat to be sure you are applying the right dose formulated for the weight of your cat. Separate the treated cat from all other cats and cats for 24 hours after treatment has been applied.

How to open: Remove product tubes from the package. Separate one tube from the others. Hold the tube with notched end pointing up and away from the face and body. Use scissors to cut off the narrow end at the notches along the lines.

How to apply: Invert tube over cat and use open end to part cat's hair. Squeeze tube firmly to apply all of the solution to the cat's skin, as directed below. Do not apply more often than once every 4 weeks.

Apply one tube as a spot on the cat's back between the shoulder blades.

Side Effects: Monitor your cat after application. Cats may experience some temporary irritation at the site of product application such as redness, scratching or other signs of discomfort. If signs of sensitivity occur, bathe your cat with a mild soap and rinse with large amounts of water. If these or other side effects occur, consult your veterinarian or call 1-844-685-9173. If your cat has an unusual reaction to the initial application, consult a veterinarian before repeating application. The most common signs of incidental ingestion are excessive salivation and foaming at the mouth. If these symptoms occur, immediately feed your cat and continue to monitor your cat for the next 24 hours.

FREQUENCY OF APPLICATION

Use ZoGuard Plus for Cats monthly for complete control of flea, tick, and chewing lice infestations. Studies show that ZoGuard Plus for Cats kills adult fleas for up to three months. Apply monthly if your cat is at high risk for flea reinfestation. Apply monthly to control ticks and chewing lice. ZoGuard Plus for Cats remains effective even after bathing, water immersion, or exposure to sunlight. Allow treated area to dry thoroughly. Do not reapply for 30 days.

If you have questions or comments about this product, please write:

Promika, LLC
1204 Village Market Pl #273, Morrisville, NC 27560

©2014 Promika. ZoGuard[™] is a trademark of Promika, LLC.
Frontline[®] is a registered trademark of Merial. ZoGuard[™] Plus for Cats is not manufactured, or distributed by Merial, seller of Frontline[®] Plus for Cats.



Distributed by:
Promika, LLC.
1204 Village Market Place #273
Morrisville, NC 27560

Contains Fipronil and S-Methoprene,
the active ingredients used in
FRONTLINE[®] Plus for Cats

**ONLY for cats & kittens
over 8 weeks of age and over 1.5 lbs**

Apply one tube (0.67 mL) (0.023 fl oz)

Three 0.017 fl oz (0.50 mL) Applicators

Net Contents: 0.051 fl oz

KILLS



FLEAS



TICKS



FLEA EGGS



FLEA LARVAE



**CHEWING
LICE**

DIRECTIONS FOR USE

How to Open

1. Remove product tubes from the package.
2. Separate one tube from the others. Hold the tube with notched end pointing up and away from the face and body. Use scissors to cut off the narrow end at the notches along the line.

How to Apply

1. Invert tube over cat and use open end to part cat's hair.
2. Squeeze tube firmly to apply all of the solution to the cat's skin as a spot on the cat's back between the shoulder blades.



STORAGE AND DISPOSAL

PESTICIDE STORAGE: Do not remove tube from the pack until ready to use. Store in a cool (below 77°F/25°C) dry place inaccessible to children and pets. Do not refrigerate. Protect from direct sunlight

CONTAINER DISPOSAL: If empty: Non-refillable container. Do not reuse or refill this container.

Offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

In Case of Emergency,
Call: 1-844-685-9173

EPA Reg. No. 53883-352-91300