

## 2(ee) Recommendation

READ AND FOLLOW THE ENTIRE LABEL BOOKLET FOR AGRIMYCIN BEFORE  
PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS PRODUCT BULLETIN



# AGRIMYCIN

EPA Reg. No. 55146-96

### Fire Blight for Pears in California

#### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

The following is a Nufarm Americas Inc. - AGT Division authorized 2(ee) recommendation. This recommendation is made as permitted by Section 2(ee) of FIFRA, as amended, and has not been submitted to or approved by the EPA.

**This 2(ee) recommendation must be in the possession of the user at the time of pesticide application.**

Fire Blight ( <b>High Disease Pressure</b> )	24-48 oz Agri-mycin per application pre acre. Equivalent to 50-100 ppm at 600 gallons per acre.	Spray trees at 20%-30% bloom.	Spray trees every 3-4 days during blossom time. Apply sprays after petal fall every 10-14 days to control twig blight. (This could mean an additional 6-8 applications after blossom sprays.) Do not apply within 30 days of harvest.
--	--	-------------------------------	---

Fire Blight	28.8 oz. Agri-Mycin 17 per acre per application (equivalent to 60 ppm at 600 gals/A)	Spray trees at 10% bloom.	Repeat at 5-day intervals until late bloom is over. This could mean 12-15 applications.) Continue to spray at 5-7 day intervals to control shoot and fruit infections. Do not apply within 30 days of harvest.
-------------	---	---------------------------	--

#### NOTICE

Read the "LIMIT of WARRANTY AND LIABILITY" in the label booklet for AgriMycin before using this product. Those terms apply to this supplemental labeling and if those terms are not acceptable, return the product unopened at once.

**Manufactured By**  
**Nufarm Americas Inc. – AGT Division**  
Burr Ridge, IL 60527