Torque™ Fungicide - Product Bulletin

EPA Reg. No. 1001-87

FIFRA Section 2ee Recommendation - For Preventative and Curative Control of Fairy Ring caused by various Basidiomycete fungi

For distribution and use only within States where Torque Fungicide is currently registered (except NY and TX).

Directions for Use

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. The following recommendation is made as permitted under FIFRA Section 2(ee)(3) and has not been submitted to or approved by the US EPA.

Torque Fungicide may be used for the Preventative and Curative Control of Fairy Ring caused by various Basidiomycete fungi.

For Preventative treatment:

Apply Torque Fungicide before fairy ring symptoms appear in spring starting when 5 day average soil temperatures at 2 inches reach 55-60F. Apply as a foliar spray at the rate of 0.6 fluid ounces per 1,000 square feet. Use sufficient water volume (66-132 GPA) and appropriate spray nozzles for through plant coverage. Water the application into the active root zone within 4 hours for best results. Only use a wetting agent as required for the penetration of active hydrophobic soil conditions. User should make a second application of Torque when the 5 day average soil temperatures reach 65-70F. Do not exceed a 28 day interval between applications. Torque may also be tank mixed or rotated with Affirm as allowed by label. Do not apply to over seeded Bermudagrass during spring transition. Complete green-up should occur before treatment to avoid potential growth reduction.

For Curative treatment:

Apply Torque Fungicide as a foliar spray at the rate of 0.6 fluid ounces per 1,000 square feet to areas where fairy ring symptoms are present. Use sufficient water volume (66-132 GPA) and appropriate spray nozzles for through plant coverage. Water the application into the active root zone within 4 hours for best results. Use a wetting agent as required for the penetration of active hydrophobic soil conditions. User may make a second application of Torque in the same manner utilizing a 28-day interval. Torque may also be tank mixed or rotated with Affirm as allowed by label.

Note: For First aid and other Precautionary statements, see label attached to the container. This recommendation is made as permitted under FIFRA Section 2(ee)(3) and has not been submitted to or approved by EPA. All applicable directions, restrictions, and precautions on the EPA registered label are to be followed. This bulletin must be in the possession of the user or applicator at the time of pesticide use.

Cleary Chemicals, LLC 178 Ridge Road, Suite A, Dayton, NJ 08810 (800) 524-1662 www.clearychemical.com

