### HERBICIDE



## TRIAD SELECT Broad spectrum weed control

ACTIVE INGREDIENTS 2,4-D – Group 4 Triclopyr – Group 4 Dicamba acid – Group 4 Sulfentrazone – Group 14

 % ACTIVE INGREDIENTS

 2,4-D
 31.82%

 Triclopyr
 8.40%

 Dicamba acid
 2.43%

 Sulfentrazone
 0.73%

FORMULATION Emulsifiable concentrate

**CHEMICAL FAMILY** PPO inhibitor (Group 14), Growth regulator (Group 4)

BRAND ALTERNATIVE T-Zone<sup>™</sup>

MODE OF ACTION Cell membrane disrupters, causes excessive growth until death.

RESTRICTED USE No

SIGNAL WORD Caution

PACKAGING 12 x 1 qt , 4 x 1 gal, 2 x 2.5 gal, 5 x 30gal

LABELED USES Residential lawns Athletic fields Golf courses Parks Industrial and institutional buildings and more



**Triad TZ Select**<sup>™</sup> contains four active ingredients, with two modes of action to provide dependable post-emergent control of broadleaf weeds in a variety of turf types. **Triad TZ Select**<sup>™</sup> is quickly taken up by the leaves and stems of plants and weed growth ceases within hours. The weed control effects may be seen within a few hours to a few days after application. Weed death occurs in 1 to 4 weeks after application, depending on the application rate, sensitivity of the weed, and environmental conditions.

### **PRODUCT BENEFITS**

- Excellent post-emergent activity with proven rapid and effective weed control performance for some of the toughest broadleaf weeds in turfgrass such as dandelion, spurge, white clover, plantains, ground ivy, Virginia buttonweed and wild violet. Labelled for the suppression of yellow nutsedge
- This product exhibits improved cool-weather performance
- · Can be applied to fully dormant bermuda, zoysia and bahia grasses
- The speed of action (rate of weed phytotoxicity, yellowing) and the early weed symptoms are features of sulfentrazone
- This product is generally rainfast in as little as three hours



Prime Source, a Division of Albaugh, LLC 1525 NE 36th Street, Ankeny, IA 50021 • www.primesource-albaugh.com

©2021 Albaugh, LLC. Prime Source and Select are trademarks of Albaugh, LLC. T-Zone is a trademark of PBI-Gordon Corporation. Always read and follow label instructions.

# TRIAD SELECT

### **APPLICATION INFORMATION**

Generally, the lower application rates within the specified range will provide satisfactory control of sensitive weed species. The higher application rates within the specified range will be required for dense infestations of perennial weeds, for adverse/extreme environmental conditions, or for weeds hardened off or more mature.

Table 1. Use rates for ornamental turfgrass, sod farms and non-cropland.						
Rate	Spray Volume					
Cool season turf						
3.25 to 4 pints/A	10 to 220 gal/A					
(1.2 to 1.5 fl. oz./1,000 sq.ft.)	(0.25 to 5.0 gal/1,000 sq.ft.)					
Warm season turf (Dormant turf)						
2 to 2.25 pints/A	10 to 220 gal/A					
(0.75 to 0.83 fl. oz./1,000 sq.ft.)	(0.25 to 5.0 gal/1,000 sq.ft.)					
	Rate         Cool season turf         3.25 to 4 pints/A         (1.2 to 1.5 fl. oz./1,000 sq.ft.)         Warm season turf (Dormant turf)         2 to 2.25 pints/A					

**Dormant turf**: This product may be applied to fully dormant bermudagrass, fully dormant zoysiagrass and fully dormant bahiagrass. **Note**: Do not apply to above listed warm-season turfgrass unless turf injury can be tolerated. It is impossible to test all environmental conditions for the listed warm-season turfgrass. We suggest testing this product on a small area and observe the treated area for 30 days to determine the acceptability of turf discoloration.

Do not apply this product to warm-season turfgrass during spring green-up or in the fall during the transition period between active growth and dormancy.

TABLE 2. Limitations on broadcast applications to ornamental turfgrass, sod farms and non-cropland.					
Use Site	Maximum rate per application	Maximum number of applications per year*	Minimum interval between applications	Maximum seasonal rate	
Ornamental turfgrass	4.0 pints/A (0.88 lb 2,4-D ae/A) (0.25 lb triclopyr ae/A) (0.10 lb dicamba ae/A) (0.03 lb sulfentrazone ae/A)	2	N/A	8.0 pints/A (1.75 lb 2,4-D ae/A) (0.50 lb triclopyr ae/A) (0.20 lb dicamba ae/A) (0.06 lb sulfentrazone ae/A)	
Sod farms	4.0 pints/A (0.88 lb 2,4-D ae/A) (0.25 lb. triclopyr ae/A) (0.10 lb dicamba ae/A) (0.03 lb sulfentrazone ae/A)	2	21 days	8.0 pints/A (1.75 lb 2,4-D ae/A) (0.50 lb triclopyr ae/A) (0.20 lb dicamba ae/A) (0.06 lb sulfentrazone ae/A)	
Non-cropland	4.0 pints/A	1 (woody plants)	N/A	8.0 pints/A (1.75 lb 2,4-D ae/A) (0.50 lb triclopyr ae/A) (0.20 lb dicamba ae/A) (0.06 lb sulfentrazone ae/A)	
	(0.88 lb 2,4-D ae/A) (0.25 lb triclopyr ae/A) (0.10 lb dicamba ae/A) (0.03 lb sulfentrazone ae/A)	2 (annual and perennial weeds)	30 days		
* New York: On	ly one application per year of th	nis product is allowed.		,	

#### Refer to label for full list of weeds controlled.